


University of Stuttgart

Department of Mathematics



Cedric
Beschle

CLMC techniques for elliptic PDEs with random discontinuities

Joint work with
Andrea Barth

30.06.23

Outline

$$u \xrightarrow{\text{adaptive FE}} u_\ell \xrightarrow{\text{q.o.i}} Q_\ell := Q(u_\ell) \xrightarrow{\text{mean}} \mathbb{E}[Q_\ell] \xrightarrow{\text{(Q)CLMC}} \hat{Q} \approx \mathbb{E}[Q_\ell]$$

Outline

$$u \xrightarrow{\text{adaptive FE}} u_\ell \xrightarrow{\text{q.o.i}} Q_\ell := \mathcal{Q}(u_\ell) \xrightarrow{\text{mean}} \mathbb{E}[Q_\ell] \xrightarrow{\text{(Q)CLMC}} \hat{Q} \approx \mathbb{E}[Q_\ell]$$

Random PDE with discontinuous coefficient and its discretization

Continuous level Monte Carlo (CLMC)


Quasi continuous level Monte Carlo (QCLMC)

Numerical experiments

Random PDE with discontinuous coefficient

Flow through fractured porous media:

- PDE with discontinuous random coefficient
- spatial **discontinuities** accounting for fractures
- randomness accounting for **uncertainties**



Simplified model problem

Define

- complete probability space $(\Omega, \mathcal{A}, \mathbb{P})$
- bounded and connected Lipschitz domain $\mathcal{D} \subset \mathbb{R}^2$
- linear random elliptic PDE: Given a and f , let u be the solution to

$$\begin{aligned} -\nabla \cdot (a(\omega, x) \nabla u(\omega, x)) &= f(x) && \text{in } \Omega \times \mathcal{D}, \\ u(\omega, x) &= 0 && \text{on } \Omega \times \partial\mathcal{D}. \end{aligned}$$

Simplified model problem

Define

- complete probability space $(\Omega, \mathcal{A}, \mathbb{P})$
- bounded and connected Lipschitz domain $\mathcal{D} \subset \mathbb{R}^2$
- linear random elliptic PDE: Given a and f , let u be the solution to

$$\begin{aligned} -\nabla \cdot (a(\omega, x) \nabla u(\omega, x)) &= f(x) && \text{in } \Omega \times \mathcal{D}, \\ u(\omega, x) &= 0 && \text{on } \Omega \times \partial\mathcal{D}. \end{aligned}$$

Coefficient $a : \Omega \times \mathcal{D} \rightarrow \mathbb{R}$:

- **random**
- **discontinuous in space**

\Rightarrow **interface problem** cf. [Babuška, 1970, Teckentrup, Scheichl, Giles and Ullmann, 2013, Barth and Stein, 2018].

Simplified model problem

Define

- complete probability space $(\Omega, \mathcal{A}, \mathbb{P})$
- bounded and connected Lipschitz domain $\mathcal{D} \subset \mathbb{R}^2$
- linear random elliptic PDE: Given a and f , let u be the solution to

$$\begin{aligned} -\nabla \cdot (a(\omega, x) \nabla u(\omega, x)) &= f(x) && \text{in } \Omega \times \mathcal{D}, \\ u(\omega, x) &= 0 && \text{on } \Omega \times \partial\mathcal{D}. \end{aligned}$$

Coefficient $a : \Omega \times \mathcal{D} \rightarrow \mathbb{R}$:

- **random**
- **discontinuous in space**

\Rightarrow **interface problem** cf. [Babuška, 1970, Teckentrup, Scheichl, Giles and Ullmann, 2013, Barth and Stein, 2018].

Remark: Weak solutions $u(\omega) \in H_0^1(\mathcal{D})$ for \mathbb{P} -almost all $\omega \in \Omega$ and


$$\|u\|_{L^p(\Omega; H_0^1)} \leq C(p, P, f, \mathcal{D}) \quad \text{for } 1 \leq p \leq \infty.$$

Sample of Box PDE coefficient and solution

$\mathcal{D} := [0, 1]^2$, for $\omega \in \Omega$, blue box:

- random center $\sim \mathcal{U}([0.4, 0.6]^2)$
- random edge length $\sim \mathcal{U}([0.2, 0.3])$
- outside $P \in \mathbb{R}_{>0}$ and inside P^{-1}

Approximation of $u(\omega)$ by $u_\ell(\omega)$ with linear finite elements on a mesh \mathcal{K}_ℓ .



Sample of Box PDE coefficient and solution


$\mathcal{D} := [0,1]^2$, for $\omega \in \Omega$, blue box:



- random center $\sim \mathcal{U}([0.4,0.6]^2)$
- random edge length $\sim \mathcal{U}([0.2,0.3])$
- outside $P \in \mathbb{R}_{>0}$ and inside P^{-1}

Approximation of $u(\omega)$ by $u_\ell(\omega)$ with linear finite elements on a mesh \mathcal{K}_ℓ .

Important: Discontinuous coefficient with large jump:

- flat areas – no refinement
- steep areas – refinement!







Unstructured uniform meshes \mathcal{K}


↓
high resolution at the discontinuity

↓
overall high resolution

↓
high computational cost



Sample adaptive meshes $\mathcal{K}(\omega)$, for $\omega \in \Omega$



Goal-oriented a-posteriori error estimator for

$$Q(u(\omega)) := \|u(\omega)\|_{H^1(\mathcal{D})},$$

[B. and Barth, '23 (1)].

Outline

Random PDE with discontinuous coefficient and its discretization

Continuous level Monte Carlo (CLMC)

Quasi continuous level Monte Carlo (QCLMC)

Numerical experiments

Continuous level Monte Carlo (CLMC)

Continuous level Monte Carlo (CLMC) [Detommaso, Dodwell and Scheichl, 2019]

- generalisation of MLMC
- continuous level of refinement $\ell \in \mathbb{R}_{\geq 0}$
- stochastic process of approximations $(Q(\ell))_{\ell \geq 0} := (\mathcal{Q}(u_\ell))_{\ell \geq 0}$
- random variable $L_r \sim \text{Exp}(r)$ with $r \in \mathbb{R}_{> 0}$
- **sample-adaptive approximations**

Continuous level Monte Carlo (CLMC)

Select

- maximal level $L_{max} \in \mathbb{R}_{>0}$
- sample number $M \in \mathbb{N}$
- $((Q^{(k)}(\ell)_{\ell \geq 0})_{k=1}^M$ i.i.d. copies of $Q(\ell)_{\ell \geq 0}$

CLMC estimator for the expectation $\mathbb{E}(Q - Q_0)$:

$$\hat{Q}_{L_{max}}^{\text{CLMC}} := \frac{1}{M} \sum_{k=1}^M \int_0^{\min\{L_{max}, L_r^{(k)}\}} \frac{1}{\mathbb{P}(L_r \geq \ell)} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) d\ell$$

Continuous level Monte Carlo (CLMC)

Select

- maximal level $L_{max} \in \mathbb{R}_{>0}$
- sample number $M \in \mathbb{N}$
- $((Q^{(k)}(\ell)_{\ell \geq 0})_{k=1}^M$ i.i.d. copies of $Q(\ell)_{\ell \geq 0}$

CLMC estimator for the expectation $\mathbb{E}(Q - Q_0)$:

$$\begin{aligned}\hat{Q}_{L_{max}}^{\text{CLMC}} &:= \frac{1}{M} \sum_{k=1}^M \int_0^{\min\{L_{max}, L_r^{(k)}\}} \frac{1}{\mathbb{P}(L_r \geq \ell)} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) d\ell \\ &= \frac{1}{M} \sum_{k=1}^M \int_0^{L_{max}} e^{r\ell} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) \mathbf{1}_{[0, L_r^{(k)}]}(\ell) d\ell.\end{aligned}$$

Continuous level Monte Carlo (CLMC)

Select

- maximal level $L_{max} \in \mathbb{R}_{>0}$
- sample number $M \in \mathbb{N}$
- $((Q^{(k)}(\ell)_{\ell \geq 0})_{k=1}^M)$ i.i.d. copies of $Q(\ell)_{\ell \geq 0}$


CLMC estimator for the expectation $\mathbb{E}(Q - Q_0)$:

$$\begin{aligned}\widehat{Q}_{L_{max}}^{\text{CLMC}} &:= \frac{1}{M} \sum_{k=1}^M \int_0^{\min\{L_{max}, L_r^{(k)}\}} \frac{1}{\mathbb{P}(L_r \geq \ell)} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) d\ell \\ &= \frac{1}{M} \sum_{k=1}^M \int_0^{L_{max}} e^{r\ell} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) \mathbf{1}_{[0, L_r^{(k)}]}(\ell) d\ell.\end{aligned}$$


Denote by $(Q_j^{(k)})_{j \geq 1}$ a sequence of approximations at level $(\ell_j^{(k)})_{j \geq 1}$. Then define via linear interpolation

$$\left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) := \frac{Q_j^{(k)} - Q_{j-1}^{(k)}}{\ell_j^{(k)} - \ell_{j-1}^{(k)}} \quad \text{for } \ell \in (\ell_{j-1}^{(k)}, \ell_j^{(k)}).$$

↑ resolution $L_r^{(k)}$




$$L_r^{(1)} = 2$$




$$L_r^{(2)} = 3$$

← M samples →



$$L_r^{(3)} = 1$$

→



$$L_r^{(M)} = 3$$

Outline

Random PDE with discontinuous coefficient and its discretization

Continuous level Monte Carlo (CLMC)

Quasi continuous level Monte Carlo (QCLMC)

Numerical experiments

Motivation: Quasi continuous level Monte Carlo

$$\hat{Q}_{L_{max}}^{\text{CLMC}} := \frac{1}{M} \sum_{k=1}^M \int_0^{L_{max}} e^{r\ell} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) \mathbf{1}_{[0, L_r^{(k)}]}(\ell) d\ell.$$

Approximation of the tail distribution

$$e^{-r\ell} = \mathbb{P}(L \geq \ell) = \mathbb{E}(\mathbf{1}_{[0, L_r]}(\ell)) \approx \frac{1}{M} \sum_{k=1}^M \mathbf{1}_{[0, L_r^{(k)}]}(\ell) \quad \text{for all } \ell \in [0, \infty).$$

\Rightarrow Generate $(L_r^{(k)}; k = 1, \dots, M)$ as a deterministic quasi random sequence.

F -discrepancy

Let $F : B \rightarrow \mathbb{R}$ be a CDF.

For a given point set $P := \{y_1, \dots, y_M\}$ the F -discrepancy is defined as

$$D_{F,P} := \sup_{y \in B} \left| \frac{1}{M} \sum_{k=1}^M \mathbb{1}_{\{y_k \leq y\}} - F(y) \right|,$$

cf. [Fang, Wang and Bentler, 1994].

F -discrepancy

Let $F : B \rightarrow \mathbb{R}$ be a CDF.

For a given point set $P := \{y_1, \dots, y_M\}$ the F -discrepancy is defined as

$$D_{F,P} := \sup_{y \in B} \left| \frac{1}{M} \sum_{k=1}^M \mathbb{1}_{\{y_k \leq y\}} - F(y) \right|,$$

cf. [Fang, Wang and Bentler, 1994].

If F is continuous with continuous inverse, then

$$D_{F,P} = D_M^*(P).$$

The same holds true for the TDF, since $T = 1 - F$.

Quasi random numbers and F -discrepancy

Lemma (B. and Barth, '23 (2))

Let $L_r \sim \text{Exp}(r)$ for some $r > 0$ and

$$L_r^{(k)} = -\frac{\ln(1 - x^{(k)})}{r}$$

from a $[0, 1)$ -uniform quasi random Sobol sequence, cf. [Sobol, 1967] and [Owen, 1998], $x^{(k)}$ for $k = 1, \dots, M$ and $M \in \mathbb{N}$. Then,

$$\sup_{x \in [0, \infty)} \left| \frac{1}{M} \sum_{m=1}^M \mathbb{1}_{[0, L_r^{(k)}]}(x) - e^{-rx} \right| \leq CM^{-1},$$

for $C > 0$ independent of M ,

Cf. [Dick and Pillichshammer, 2014] for the convergence rate.

F -discrepancy convergence, for $L_r = \text{Exp}(1.3)$

$$F\text{-discrepancy: } \sup_{\ell \in \mathbb{R}_+} \left| \frac{1}{M} \sum_{k=1}^M \mathbf{1}_{[0, L_r^{(k)}]}(\ell) - e^{-r\ell} \right|$$




Illustration of tail approximation







Illustration of tail approximation







Illustration of tail approximation






Illustration of tail approximation







Illustration of tail approximation







Illustration of tail approximation











Illustration of tail approximation



Remark:
Exp(r) in (Q)CLMC
is always
one-dimensional.



No additional cost to
generate the quasi
random numbers.

Quasi continuous level Monte Carlo (QCLMC)

Recently developed in [B. and Barth, '23 (1)] and analysed in [B. and Barth, '23 (2)].
Select

- deterministic quasi random sequence $L_r^{(k)}$ mimicing an exponential distribution $\text{Exp}(r)$, $r > 0$
- maximal level $L_{max} := \min\{\max_{k \in (1, \dots, M)} L_r^{(k)}, L\}$, where $L \in \mathbb{R}_{>0}$

QCLMC estimator for the expectation $\mathbb{E}(Q - Q_0)$:

$$\hat{Q}_{L_{max}}^{\text{QCLMC}} := \frac{1}{M} \sum_{k=1}^M \int_0^{L_{max}} e^{r\ell} \left(\frac{dQ}{d\ell} \right)^{(k)}(\ell) \mathbb{1}_{[0, L_r^{(k)}]}(\ell) d\ell.$$

Complexity Theorem (QCLMC)

Theorem (B. and Barth, '23 (2))

Suppose there exist positive constants $\alpha, \beta, \gamma, c_1, c_2, c_3$, with $\beta < 4\alpha$, such that for any $\ell > 0$:

$$\mathbb{E} \left[\frac{dQ}{d\ell} \right] \leq c_1 e^{-\alpha\ell} \quad \text{and} \quad \mathbb{V} \left[\frac{dQ}{d\ell} \right] \leq c_2 e^{-\beta\ell} \quad \text{and} \quad \mathcal{C}(\ell) \leq c_3 e^{\gamma\ell}.$$

Complexity Theorem (QCLMC)

Theorem (B. and Barth, '23 (2))

Suppose there exist positive constants $\alpha, \beta, \gamma, c_1, c_2, c_3$, with $\beta < 4\alpha$, such that for any $\ell > 0$:

$$\mathbb{E} \left[\frac{dQ}{d\ell} \right] \leq c_1 e^{-\alpha\ell} \quad \text{and} \quad \mathbb{V} \left[\frac{dQ}{d\ell} \right] \leq c_2 e^{-\beta\ell} \quad \text{and} \quad \mathcal{C}(\ell) \leq c_3 e^{\gamma\ell}.$$

Further, let $(L_r^{(k)}; k = 1, \dots, M)$ be a deterministic quasi random sequence obtained by inverse transformation, and let $r \in [\min\{\beta, 2\alpha, \gamma\}, \max\{\min\{\beta, 2\alpha\}, \gamma\}]$.

Complexity Theorem (QCLMC)

Theorem (B. and Barth, '23 (2))

Suppose there exist positive constants $\alpha, \beta, \gamma, c_1, c_2, c_3$, with $\beta < 4\alpha$, such that for any $\ell > 0$:

$$\mathbb{E} \left[\frac{dQ}{d\ell} \right] \leq c_1 e^{-\alpha\ell} \quad \text{and} \quad \mathbb{V} \left[\frac{dQ}{d\ell} \right] \leq c_2 e^{-\beta\ell} \quad \text{and} \quad \mathcal{C}(\ell) \leq c_3 e^{\gamma\ell}.$$

Further, let $(L_r^{(k)}; k = 1, \dots, M)$ be a deterministic quasi random sequence obtained by inverse transformation, and let $r \in [\min\{\beta, 2\alpha, \gamma\}, \max\{\min\{\beta, 2\alpha\}, \gamma\}]$. Then, there exist $L \in \mathbb{R}_{>0}$ and $M \in \mathbb{N}$, such that for any $\varepsilon \in (0, \frac{1}{e})$ and $\bar{C} > 0$ independent of M and ε

$$\text{MSE}^{\text{QCLMC}} \leq \varepsilon^2 \quad \text{and} \quad \mathcal{C}[\hat{Q}_{L_{max}}^{\text{QCLMC}}] \leq \bar{C} \varepsilon^{-2 - \max\{0, \frac{\gamma - \min\{\beta, 2\alpha\}}{\alpha}\}} |\ln(\varepsilon)|^{\delta_{r, \min\{\beta, 2\alpha\}} + \delta_{r, \gamma}},$$

where $L_{max} = \min\{L, \max_{k \in \{1, \dots, M\}} L_r^{(k)}\}$ and δ is the Dirac function.

Outline

Random PDE with discontinuous coefficient and its discretization

Continuous level Monte Carlo (CLMC)

Quasi continuous level Monte Carlo (QCLMC)

Numerical experiments

Convergence experiments

We choose

$$Q(u(\omega)) := \|u(\omega)\|_{H^1(\mathcal{D})}.$$

Comparison - time to MSE performance:

- CLMC on sample adaptive meshes, $\alpha \approx 1$, $\beta \approx 2.3$, $\gamma \approx 1$
- QCLMC on sample adaptive meshes
- MLMC on unstructured uniform meshes, $\alpha \approx 0.5$, $\beta \approx 1.4$, $\gamma \approx 1$

[Barth, Schwab and Zollinger, 2011]


[Cliffe, Giles, Scheichl and Teckentrup, 2011]

[Teckentrup, Scheichl, Giles and Ullmann, 2013]

[Haji-Ali, Nobile, von Schwerin and Tempone, 2016]

Time to MSE performance - box coefficient

$P = 300$





MSE: $\varepsilon^2 \in \{0.04, 0.025, 0.016, 0.01, 0.007, 0.004, 0.003\}$, [B. and Barth, '23 (1)].

Time to MSE performance - box coefficient

$P = 300$

$P = 1000$



MSE: $\varepsilon^2 \in \{0.04, 0.025, 0.016, 0.01, 0.007, 0.004, 0.003\}$, [B. and Barth, '23 (1)].

Summary and conclusion

Random discontinuous coefficient in PDE: negative effect on

- regularity of pathwise weak solution
- pathwise convergence rate on standard meshes (MLMC)

⇒ better use adaptive meshes (CLMC)

New QCLMC estimator

- much better tail estimate for $L_r \sim \text{Exp}(r)$
- variance reduction

Summary and conclusion

Numerical experiments:

- CLMC should outperform MLMC as solution samples have distinct areas with high error contributions
- pseudo random numbers for sampling $L_r \sim \text{Exp}(r)$ renders it worse than MLMC

⇒ QCLMC outperforms both methods

Outlook:

- apply QCLMC to random Burgers equation (discontinuous solutions)
- develop improved automatic stopping criterion for QCLMC

Preprints of submitted work

- (1) C. A. Beschle, A. Barth: Quasi Continuous Level Monte Carlo for Random Elliptic PDEs, in MCQMC 2022 Proc., submitted 2023,
<https://doi.org/10.48550/arXiv.2303.08694>.
- (2) C. A. Beschle, A. Barth: Quasi Continuous Level Monte Carlo, in ESAIM M2AN - Special issue - To commemorate Assyr Abdulle, submitted 2023,
<https://doi.org/10.48550/arXiv.2305.15949>.

Literature 1

- Barth, A., Schwab, Ch. and Zollinger, N., Multi-level Monte Carlo Finite Element method for elliptic PDEs with stochastic coefficients, Numer. Math., [2011]
- Teckentrup, A. L., Scheichl, R., Giles, M. B. and Ullmann, E., Further analysis of multilevel Monte Carlo methods for elliptic PDEs with random coefficients, Numer. Math., [2013]
- Cliffe, K., Giles, M. B., Scheichl R. and Teckentrup, A. L., Multilevel Monte Carlo methods and applications to elliptic PDEs with random coefficients, Comput. Vis. Sci., [2011]
- Babuška, I., The finite element method for elliptic equations with discontinuous coefficients, Computing (Arch. Elektron. Rechnen, [1970]
- Barth, A. and Stein, A., A study of elliptic partial differential equations with jump diffusion coefficients, SIAM/ASA J. Uncertain. Quantif., [2018]
- Detommaso, G., Dodwell, T. and Scheichl, R., Continuous level Monte Carlo and sample-adaptive model hierarchies, SIAM/ASA J. Uncertain. Quantif., [2019]
- Dick, J. and Pillichshammer F., Discrepancy theory and quasi-Monte Carlo integration, A panorama of discrepancy theory, volume 2107 of Lecture Notes in Math., [2014]

Literature 2

- Haji-Ali, A. and Nobile, F. and von Schwerin, E. and Tempone, R., Optimization of mesh hierarchies in multilevel Monte Carlo samplers, *Stoch. Partial Differ. Equ. Anal. Comput.*, [2016]
- Sobol, I.M., Distribution of points in a cube and approximate evaluation of integrals, *U.S.S.R Comput. Maths. Math. Phys.* [1967]
- Owen, A.B., Scrambling Sobol' and Niederreiter–Xing points, *J. Complexity*, [1998]
- Fang, K. T. and Wang, Y. and Bentler, P. M., Some applications of number-theoretic methods in statistics, *Statist. Sci.* [1994]