


Multilevel Monte Carlo Methods for Parametric Expectations: Distribution and Robustness Measures


Sebastian Krumscheid (sebastian.krumscheid@kit.edu) | June 29, 2023

joint work with: Q. Ayoul-Guilmar, S. Ganesh, and F. Nobile


Forward propagation of Uncertainties




- Random input parameters ξ , possibly infinite dimensional
- **Model:** for ξ given, find u s.t. $\mathcal{M}_\xi(u) = 0 \rightsquigarrow$ **Quantity of interest:** $Q(\xi) = Q(u(\xi))$



Forward propagation of Uncertainties




- Random input parameters ξ , possibly infinite dimensional
- **Model:** for ξ given, find u s.t. $\mathcal{M}_\xi(u) = 0 \rightsquigarrow$ **Quantity of interest:** $Q(\xi) = Q(u(\xi))$




- **Goal:** compute statistical quantities of Q with minimal regularity assumptions.

Forward propagation of Uncertainties




- Random input parameters ξ , possibly infinite dimensional
- **Model:** for ξ given, find u s.t. $\mathcal{M}_\xi(u) = 0 \rightsquigarrow$ **Quantity of interest:** $Q(\xi) = Q(u(\xi))$



- **Goal:** compute statistical quantities of Q with minimal regularity assumptions.


Multilevel Monte Carlo for expected values

- **Main idea:** use a hierarchy of $L \in \mathbb{N}_0$ discretizations $h_0 > h_1 > \dots > h_L$, instead of just one fine level:



Multilevel Monte Carlo for expected values

- **Main idea:** use a hierarchy of $L \in \mathbb{N}_0$ discretizations $h_0 > h_1 > \dots > h_L$, instead of just one fine level:




- The multilevel Monte Carlo (MLMC) estimator of $\mathbb{E}(Q)$ then is ($Q_\ell \equiv Q_{h_\ell}$):

$$\mathbb{E}(Q) \approx \mathbb{E}(Q_L) = \mathbb{E}(Q_0) + \sum_{\ell=1}^L \mathbb{E}(Q_\ell - Q_{\ell-1}) \approx E_{N_0}(Q_0) + \sum_{\ell=1}^L E_{N_\ell}(Q_\ell - Q_{\ell-1}) =: \hat{Q}_{N,L}$$

Multilevel Monte Carlo for expected values

- **Main idea:** use a hierarchy of $L \in \mathbb{N}_0$ discretizations $h_0 > h_1 > \dots > h_L$, instead of just one fine level:



- The multilevel Monte Carlo (MLMC) estimator of $\mathbb{E}(Q)$ then is ($Q_\ell \equiv Q_{h_\ell}$):

$$\mathbb{E}(Q) \approx \mathbb{E}(Q_L) = \mathbb{E}(Q_0) + \sum_{\ell=1}^L \mathbb{E}(Q_\ell - Q_{\ell-1}) \approx E_{N_0}(Q_0) + \sum_{\ell=1}^L E_{N_\ell}(Q_\ell - Q_{\ell-1}) =: \hat{Q}_{N,L}$$

- **MSE accuracy** of multilevel Monte Carlo method is

$$\text{MSE} \equiv \mathbb{E}\left(|\mathbb{E}(Q) - \hat{Q}_{N,L}|^2\right) = \underbrace{(\mathbb{E}(Q) - \mathbb{E}(Q_L))^2}_{\text{squared bias}} + \underbrace{\sum_{\ell=0}^L \frac{\text{Var}(Q_\ell - Q_{\ell-1})}{N_\ell}}_{\text{statistical error}}, \quad Q_{-1} \equiv 0$$

- Select parameters $L \in \mathbb{N}_0$ and $N \equiv (N_0, N_1, \dots, N_L)^T \in \mathbb{N}^{L+1}$ s.t. **MSE criterion is met at minimal cost.**
- Optimal parameters by **balancing errors**:
 - **bias = discretization error** $\rightsquigarrow L$,
 - **determine** N_ℓ by minimizing $\sum_{\ell=1}^L N_\ell C_\ell$ subject to $\text{MSE} \leq \varepsilon^2$,
 - where $C_\ell = \text{cost for each } (Q_\ell^{(i,\ell)}, Q_{\ell-1}^{(i,\ell)})$.

- Select parameters $L \in \mathbb{N}_0$ and $N \equiv (N_0, N_1, \dots, N_L)^T \in \mathbb{N}^{L+1}$ s.t. **MSE criterion is met at minimal cost.**
- Optimal parameters by **balancing errors**:
 - **bias = discretization error** $\rightsquigarrow L$,
 - **determine** N_ℓ by minimizing $\sum_{\ell=1}^L N_\ell C_\ell$ subject to **MSE** $\leq \epsilon^2$,
 - where $C_\ell = \text{cost for each } (Q_\ell^{(i,\ell)}, Q_{\ell-1}^{(i,\ell)})$.

Theorem: Complexity analysis MLMC for $h_{\ell-1}/h_\ell = s > 1$

[Giles, 2008; Cliffe et al., 2011]

- Suppose that:
- $|\mathbb{E}(Q - Q_\ell)| = \mathcal{O}(h_\ell^\alpha)$,
 - $\text{Var}(Q_\ell - Q_{\ell-1}) = \mathcal{O}(h_\ell^\beta)$,
 - $C_\ell = \mathcal{O}(h_\ell^{-\gamma})$.



If $2\alpha \geq \min\{\beta, \gamma\}$, then there exists an MLMC estimator $\hat{Q}_{N,L}$ that satisfies **MSE** $\leq \epsilon^2$ with

$$\text{comp. cost MLMC} = \sum_{\ell=1}^L N_\ell C_\ell \lesssim \begin{cases} \epsilon^{-2}, & \beta > \gamma, \\ \epsilon^{-2} \ln(\epsilon)^2, & \beta = \gamma, \\ \epsilon^{-(2 + \frac{\gamma - \beta}{\alpha})}, & \beta < \gamma. \end{cases}$$

- Other error criteria are possible, e.g., tuning MLMC for *probability of failure* is possible thanks to CLT [Collier et al. 2015] and [Hoel, K. 2019].



Assessing the distribution beyond moments

- Expectation is a **good starting point**, but we would like more: **assessing the distribution**



Assessing the distribution beyond moments

- Expectation is a **good starting point**, but we would like more: **assessing the distribution**





- One could estimate **central moments** of arbitrary order

$$\mu_p(Q) := \mathbb{E}[(Q - \mathbb{E}(Q))^p], \quad p \in \mathbb{N}$$

with MLMC [Bierig, Chernov. 2016] [K., Nobile, and Pisaroni. 2020]

Assessing the distribution beyond moments

- Expectation is a **good starting point**, but we would like more: **assessing the distribution**



- One could estimate **central moments** of arbitrary order

$$\mu_p(Q) := \mathbb{E}[(Q - \mathbb{E}(Q))^p], \quad p \in \mathbb{N}$$

with MLMC [Bierig, Chernov. 2016] [K., Nobile, and Pisaroni. 2020]

- However, some applications require techniques that systematically go **beyond using a few moments**: characterization of entire **distribution** or applications in **risk averse optimization**, e.g., involving **quantiles**.

Parametric Expectations

- Many such distributional assessments are related to **parametric expectations** of the form

$$\Phi(\vartheta) := \mathbb{E}(\phi(\vartheta, Q)) , \quad \vartheta \in \Theta$$

Parametric Expectations

- Many such distributional assessments are related to **parametric expectations** of the form

$$\Phi(\vartheta) := \mathbb{E}(\phi(\vartheta, Q)) , \quad \vartheta \in \Theta$$

Examples:

1. **Mean & variance:** $\phi(\vartheta, Q) = (Q - \vartheta)^2 \rightsquigarrow \text{Var}(Q) = \min_{\vartheta} \Phi(\vartheta), \mathbb{E}(Q) = \arg \min_{\vartheta} \Phi(\vartheta)$
1. **Characteristic function** of Q : $\phi(\vartheta, Q) = e^{i\vartheta Q} = \cos(\vartheta Q) + i \sin(\vartheta Q)$
2. **Cumulative distribution function** of Q : $\phi(\vartheta, Q) = I(Q \leq \vartheta)$

Parametric Expectations

- Many such distributional assessments are related to **parametric expectations** of the form

$$\Phi(\vartheta) := \mathbb{E}(\phi(\vartheta, Q)) , \quad \vartheta \in \Theta$$

Examples:

- Mean & variance:** $\phi(\vartheta, Q) = (Q - \vartheta)^2 \rightsquigarrow \text{Var}(Q) = \min_{\vartheta} \Phi(\vartheta), \mathbb{E}(Q) = \arg \min_{\vartheta} \Phi(\vartheta)$
- Characteristic function of Q :** $\phi(\vartheta, Q) = e^{i\vartheta Q} = \cos(\vartheta Q) + i \sin(\vartheta Q)$
- Cumulative distribution function of Q :** $\phi(\vartheta, Q) = I(Q \leq \vartheta)$

Difficulty for CDF: $\phi(\vartheta, \cdot)$ is **discontinuous**! Consequently, the variance $\text{Var}(\phi(\theta_j, Q_\ell) - \phi(\theta_j, Q_{\ell-1}))$ will decay slowly \rightsquigarrow **Not much gain in MLMC.**

Parametric Expectations

- Many such distributional assessments are related to **parametric expectations** of the form

$$\Phi(\vartheta) := \mathbb{E}(\phi(\vartheta, Q)) , \quad \vartheta \in \Theta$$

Examples:

- Mean & variance:** $\phi(\vartheta, Q) = (Q - \vartheta)^2 \rightsquigarrow \text{Var}(Q) = \min_{\vartheta} \Phi(\vartheta), \mathbb{E}(Q) = \arg \min_{\vartheta} \Phi(\vartheta)$
- Characteristic function of Q :** $\phi(\vartheta, Q) = e^{i\vartheta Q} = \cos(\vartheta Q) + i \sin(\vartheta Q)$
- Cumulative distribution function of Q :** $\phi(\vartheta, Q) = I(Q \leq \vartheta)$

Difficulty for CDF: $\phi(\vartheta, \cdot)$ is **discontinuous**! Consequently, the variance $\text{Var}(\phi(\theta_j, Q_\ell) - \phi(\theta_j, Q_{\ell-1}))$ will decay slowly \rightsquigarrow **Not much gain in MLMC.**

Remedies in the context of MLMC (cf. [Giles. 2023]):

- Regularizing ϕ to yield $\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q)) \approx \mathbb{E}(\phi_\delta(\vartheta, Q))$ [Giles, Nagapetyan, Ritter. 2015]
- Approximate CDF (PDF) based on MLMC estimates of moments [Bierig, Chernov. 2016]
- numerical smoothing via conditioning [Bayer, Ben Hammouda, Tempone. 2022]

Remedy for CDF: antiderivative/integration approach [K., Nobile. 2018]

For any $\tau \in (0, 1)$ define

$$\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q)) , \quad \phi(\vartheta, Q) = \vartheta + \frac{1}{1-\tau}(Q - \vartheta)^+ .$$

Then

$$F(\vartheta) = (1 - \tau)\Phi'(\vartheta) + \tau ,$$

which is the **starting point for the MLMC estimator**:

\rightsquigarrow requires effective estimator of Φ and its derivatives.

Remedy for CDF: antiderivative/integration approach [K., Nobile. 2018]

For any $\tau \in (0, 1)$ define

$$\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q)) , \quad \phi(\vartheta, Q) = \vartheta + \frac{1}{1-\tau}(Q - \vartheta)^+ .$$

Then

$$F(\vartheta) = (1 - \tau)\Phi'(\vartheta) + \tau ,$$

which is the **starting point for the MLMC estimator**:

\rightsquigarrow requires effective estimator of Φ and its derivatives.

This approach may also offer approximations for:

- PDF: $f(\vartheta) = F'(\vartheta) = (1 - \tau)\Phi''(\vartheta)$
- τ -quantile: $q_\tau = \inf\{\vartheta: F(\vartheta) \geq \tau\} = \arg \min_{\vartheta \in \mathbb{R}} \Phi(\vartheta)$
- Conditional Value at Risk: $\text{CVaR}_\tau = \frac{1}{1-\tau} \int_{q_\tau}^\infty \vartheta dF(\vartheta) = \min_{\vartheta \in \mathbb{R}} \Phi(\vartheta)$

Remedy for CDF: antiderivative/integration approach [K., Nobile. 2018]

For any $\tau \in (0, 1)$ define

$$\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q)) , \quad \phi(\vartheta, Q) = \vartheta + \frac{1}{1-\tau}(Q - \vartheta)^+ .$$

Then

$$F(\vartheta) = (1 - \tau)\Phi'(\vartheta) + \tau ,$$

which is the **starting point for the MLMC estimator**:

\rightsquigarrow requires effective estimator of Φ and its derivatives.

This approach may also offer approximations for:

- PDF: $f(\vartheta) = F'(\vartheta) = (1 - \tau)\Phi''(\vartheta)$
- τ -quantile: $q_\tau = \inf\{\vartheta: F(\vartheta) \geq \tau\} = \arg \min_{\vartheta \in \mathbb{R}} \Phi(\vartheta)$
- Conditional Value at Risk: $\text{CVaR}_\tau = \frac{1}{1-\tau} \int_{q_\tau}^\infty \vartheta dF(\vartheta) = \min_{\vartheta \in \mathbb{R}} \Phi(\vartheta) \rightsquigarrow \text{Fabio's talk}$

MLMC method for parametric expectations


- **Goal:** given ϕ , approximate $\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q))$ and its derivatives uniformly in Θ .

MLMC method for parametric expectations

- **Goal:** given ϕ , approximate $\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q))$ and its derivatives uniformly in Θ .
- **Idea:** extend collection of pointwise estimators in nodes $\theta = (\theta_1, \dots, \theta_n)$ to function:

MLMC method for parametric expectations

- **Goal:** given ϕ , approximate $\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q))$ and its derivatives **uniformly** in Θ .
- **Idea:** extend collection of pointwise estimators in nodes $\theta = (\theta_1, \dots, \theta_n)$ to function:




Interpolation approach (e.g., splines or polynomials):

- introduce grid $\theta = (\theta_1, \dots, \theta_n) \subset \Theta$
- compute MLMC estimate $\Phi_L^{\text{MLMC}}(\theta_j)$ of $\Phi(\theta_j)$, $j = 1, \dots, n$ (same samples of Q_ℓ for every θ_j)
- interpolate values $\Phi_L^{\text{MLMC}}(\theta) = (\Phi_L^{\text{MLMC}}(\theta_j))_{j=1}^n$:
 $\hat{\Phi}_L = \mathcal{I}_n(\Phi_L^{\text{MLMC}}(\theta))$

MLMC method for parametric expectations

- **Goal:** given ϕ , approximate $\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q))$ and its derivatives **uniformly** in Θ .
- **Idea:** extend collection of pointwise estimators in nodes $\theta = (\theta_1, \dots, \theta_n)$ to function:



Interpolation approach (e.g., splines or polynomials):

- introduce grid $\theta = (\theta_1, \dots, \theta_n) \subset \Theta$
- compute MLMC estimate $\Phi_L^{\text{MLMC}}(\theta_j)$ of $\Phi(\theta_j)$, $j = 1, \dots, n$ (same samples of Q_ℓ for every θ_j)
- interpolate values $\Phi_L^{\text{MLMC}}(\theta) = (\Phi_L^{\text{MLMC}}(\theta_j))_{j=1}^n$:
 $\hat{\Phi}_L = \mathcal{I}_n(\Phi_L^{\text{MLMC}}(\theta))$

Exemplary **a-priori properties** for **Spline interpolation** \mathcal{I}_n :

- $\left\| f^{(m)} - \frac{d^m}{d\vartheta^m} \mathcal{I}_n(f(\theta)) \right\|_{L^\infty} \leq c_1 n^{m-(k+1)}$, for any $f \in C^{k+1}(\bar{\Theta})$, $m \leq k$
- $\left\| \mathcal{I}_n(\mathbf{x}) \right\|_{L^\infty} \leq c_2 \|\mathbf{x}\|_{\ell^\infty}$, for any $\mathbf{x} \in \mathbb{R}^n$
- $\text{cost}(\mathcal{I}_n(\mathbf{x})) \leq c_3 n$, $\mathbf{x} \in \mathbb{R}^n$
- $\left\| \frac{d^m}{d\vartheta^m} \mathcal{I}_n(\mathbf{x}) \right\|_{L^\infty} \leq c_5 (n-1)^m \left\| \mathcal{I}_n(\mathbf{x}) \right\|_{L^\infty}$, $\mathbf{x} \in \mathbb{R}^n$, $m \geq 1$ (**inverse inequality**)

Complexity Analysis

- Measure accuracy in terms of mean squared error:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) := \mathbb{E} \left(\left\| \hat{\Phi}_L^{(m)} - \Phi^{(m)} \right\|_{L^\infty}^2 \right), \quad m \in \mathbb{N}_0, \quad L^\infty \equiv L^\infty(\Theta).$$

Complexity Analysis

- Measure accuracy in terms of mean squared error:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) := \mathbb{E} \left(\left\| \hat{\Phi}_L^{(m)} - \Phi^{(m)} \right\|_{L^\infty}^2 \right), \quad m \in \mathbb{N}_0, \quad L^\infty \equiv L^\infty(\Theta).$$

- Using the notation $\Phi_L = \mathbb{E}(\phi(\cdot, Q_L))$, the MSE can be decomposed as:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) \leq \underbrace{3 \left\| \Phi^{(m)} - \mathcal{I}_n^{(m)}(\Phi) \right\|_{L^\infty}^2}_{\text{squared interpolation error}} + \underbrace{3 \left\| \mathcal{I}_n^{(m)}(\Phi - \Phi_L) \right\|_{L^\infty}^2}_{\text{squared bias}} + \underbrace{3 \mathbb{E} \left(\left\| \mathcal{I}_n^{(m)}(\Phi_L - \Phi_L^{\text{MLMC}}) \right\|_{L^\infty}^2 \right)}_{\text{stat. error}}$$

Complexity Analysis

- Measure accuracy in terms of mean squared error:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) := \mathbb{E} \left(\left\| \hat{\Phi}_L^{(m)} - \Phi^{(m)} \right\|_{L^\infty}^2 \right), \quad m \in \mathbb{N}_0, \quad L^\infty \equiv L^\infty(\Theta).$$

- Using the notation $\Phi_L = \mathbb{E}(\phi(\cdot, Q_L))$, the MSE can be decomposed as:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) \leq \underbrace{3 \left\| \Phi^{(m)} - \mathcal{I}_n^{(m)}(\Phi) \right\|_{L^\infty}^2}_{\text{squared interpolation error}} + \underbrace{3 \left\| \mathcal{I}_n^{(m)}(\Phi - \Phi_L) \right\|_{L^\infty}^2}_{\text{squared bias}} + \underbrace{3 \mathbb{E} \left(\left\| \mathcal{I}_n^{(m)}(\Phi_L - \Phi_L^{\text{MLMC}}) \right\|_{L^\infty}^2 \right)}_{\text{stat. error}}$$

- Parameter selection:**

- interpolation error $\rightsquigarrow n$,
- bias $\rightsquigarrow L$,
- cost minimization s.t. **variance constraint** $\rightsquigarrow N_\ell$

Complexity Analysis

- Measure accuracy in terms of mean squared error:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) := \mathbb{E} \left(\left\| \hat{\Phi}_L^{(m)} - \Phi^{(m)} \right\|_{L^\infty}^2 \right), \quad m \in \mathbb{N}_0, \quad L^\infty \equiv L^\infty(\Theta).$$

- Using the notation $\Phi_L = \mathbb{E}(\phi(\cdot, Q_L))$, the MSE can be decomposed as:

$$\text{MSE}(\hat{\Phi}_L^{(m)}) \leq \underbrace{3 \left\| \Phi^{(m)} - \mathcal{I}_n^{(m)}(\Phi) \right\|_{L^\infty}^2}_{\text{squared interpolation error}} + \underbrace{3 \left\| \mathcal{I}_n^{(m)}(\Phi - \Phi_L) \right\|_{L^\infty}^2}_{\text{squared bias}} + \underbrace{3 \mathbb{E} \left(\left\| \mathcal{I}_n^{(m)}(\Phi_L - \Phi_L^{\text{MLMC}}) \right\|_{L^\infty}^2 \right)}_{\text{stat. error}}$$

- Parameter selection:**

- interpolation error $\rightsquigarrow n$,
- bias $\rightsquigarrow L$,
- cost minimization s.t. **variance constraint** $\rightsquigarrow N_\ell$

- Notation:** $\text{Var}(\xi) = \mathbb{E}(\|\xi - \mathbb{E}(\xi)\|_{\ell^\infty}^2)$, for any r.v. ξ with values in \mathbb{R}^n

- Useful technical result:** let $(\xi^{(1)}, \dots, \xi^{(N)}) \subset \mathbb{R}^n$ independent, then

$$\text{Var} \left(\sum_{i=1}^N \xi^{(i)} \right) \leq c \ln(n) \sum_{i=1}^N \text{Var}(\xi^{(i)})$$

Theorem: A-priori MSE complexity ($h_{\ell-1}/h_\ell = s > 1$) for Spline interpolation [K., Nobile. 2018]

- Suppose that:
- $\sup_{\vartheta \in \Theta} |\Phi(\vartheta) - \mathbb{E}(\phi(\vartheta, \mathbf{Q}_l))| = \mathcal{O}(h_l^\alpha)$
 - $\mathbb{E}\left(\|\phi(\cdot, \mathbf{Q}_l) - \phi(\cdot, \mathbf{Q}_{l-1})\|_{L^\infty(\Theta)}^2\right) = \mathcal{O}(h_l^\beta),$
 - cost for each $(\mathbf{Q}_\ell^{(i,\ell)}, \mathbf{Q}_{\ell-1}^{(i,\ell)}) = \mathcal{O}(h_\ell^{-\gamma})$

Let $m = 0$. If $\Phi \in C^{k+1}(\Theta)$ and $2\alpha \geq \min\{\beta, \gamma\}$, then there exists an MLMC estimator $\hat{\Phi}_L$ of Φ such that $\text{MSE}(\hat{\Phi}_L) = \mathcal{O}(\varepsilon^2)$ with

$$\text{comp. cost} \lesssim \varepsilon^{-(2+\frac{1}{k+1})} |\ln(\varepsilon)| + |\ln(\varepsilon)| \begin{cases} \varepsilon^{-2}, & \beta > \gamma, \\ \varepsilon^{-2} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2+\frac{\gamma-\beta}{\alpha})}, & \beta < \gamma. \end{cases}$$

- **NB:** first term accounts for cost of computing interpolation: negligible for heavy computational models; systematically removable by $n = n_\ell$ (different interpolation grid on each level; Cor. 2.3 in [K., Nobile. 2018]).

Theorem: A-priori MSE complexity ($h_{\ell-1}/h_\ell = s > 1$) for Spline interpolation [K., Nobile. 2018]

- Suppose that:
- $\sup_{\vartheta \in \Theta} |\Phi(\vartheta) - \mathbb{E}(\phi(\vartheta, \mathbf{Q}_l))| = \mathcal{O}(h_l^\alpha)$
 - $\mathbb{E}\left(\|\phi(\cdot, \mathbf{Q}_l) - \phi(\cdot, \mathbf{Q}_{l-1})\|_{L^\infty(\Theta)}^2\right) = \mathcal{O}(h_l^\beta),$
 - cost for each $(\mathbf{Q}_\ell^{(i,\ell)}, \mathbf{Q}_{\ell-1}^{(i,\ell)}) = \mathcal{O}(h_\ell^{-\gamma})$

Let $m = 0$. If $\Phi \in C^{k+1}(\Theta)$ and $2\alpha \geq \min\{\beta, \gamma\}$, then there exists an MLMC estimator $\hat{\Phi}_L$ of Φ such that $\text{MSE}(\hat{\Phi}_L) = \mathcal{O}(\varepsilon^2)$ with

$$\text{comp. cost} \lesssim \varepsilon^{-(2+\frac{1}{k+1})} |\ln(\varepsilon)| + |\ln(\varepsilon)| \begin{cases} \varepsilon^{-2}, & \beta > \gamma, \\ \varepsilon^{-2} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2+\frac{\gamma-\beta}{\alpha})}, & \beta < \gamma. \end{cases}$$

- **NB:** first term accounts for cost of computing interpolation: negligible for heavy computational models; systematically removable by $n = n_\ell$ (different interpolation grid on each level; Cor. 2.3 in [K., Nobile. 2018]).
- Neglecting first term: complexity is the same as for expectations, up to extra log factor.

Theorem: A-priori MSE complexity ($h_{\ell-1}/h_\ell = s > 1$) for Spline interpolation [K., Nobile. 2018]

- Suppose that:
- $\sup_{\vartheta \in \Theta} |\Phi(\vartheta) - \mathbb{E}(\phi(\vartheta, \mathbf{Q}_l))| = \mathcal{O}(h_l^\alpha)$
 - $\mathbb{E}(\|\phi(\cdot, \mathbf{Q}_l) - \phi(\cdot, \mathbf{Q}_{l-1})\|_{L^\infty(\Theta)}^2) = \mathcal{O}(h_l^\beta),$
 - cost for each $(\mathbf{Q}_\ell^{(i,\ell)}, \mathbf{Q}_{\ell-1}^{(i,\ell)}) = \mathcal{O}(h_\ell^{-\gamma})$

Let $m = 0$. If $\Phi \in C^{k+1}(\Theta)$ and $2\alpha \geq \min\{\beta, \gamma\}$, then there exists an MLMC estimator $\hat{\Phi}_L$ of Φ such that $\text{MSE}(\hat{\Phi}_L) = \mathcal{O}(\varepsilon^2)$ with

$$\text{comp. cost} \lesssim \varepsilon^{-(2+\frac{1}{k+1})} |\ln(\varepsilon)| + |\ln(\varepsilon)| \begin{cases} \varepsilon^{-2}, & \beta > \gamma, \\ \varepsilon^{-2} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2+\frac{\gamma-\beta}{\alpha})}, & \beta < \gamma. \end{cases}$$

- **NB:** first term accounts for cost of computing interpolation: negligible for heavy computational models; systematically removable by $n = n_\ell$ (different interpolation grid on each level; Cor. 2.3 in [K., Nobile. 2018]).
- Neglecting first term: complexity is the same as for expectations, up to extra log factor.
- Similar complexity for Φ analytic and using an interpolation in global polynomials:

$$\text{comp. cost} \lesssim \varepsilon^{-2} |\ln(\varepsilon)|^4 + |\ln(\varepsilon)|^3 \begin{cases} \varepsilon^{-2}, & \beta > \gamma, \\ \varepsilon^{-2} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2+\frac{\gamma-\beta}{\alpha})} |\ln(\varepsilon)|^{\frac{\gamma-\beta}{\alpha}}, & \beta < \gamma. \end{cases}$$

Theorem (cont.)

If $\Phi \in C^{k+1}(\Theta)$ and $m \leq k$, then there exists $\hat{\Phi}_L$ such that $\text{MSE}(\hat{\Phi}_L^{(m)}) = \mathcal{O}(\varepsilon^2)$:

$$\text{comp. cost (no interp. cost)} \lesssim |\ln(\varepsilon)| \begin{cases} \varepsilon^{-2 \frac{k+1}{k+1-m}}, & \beta > \gamma, \\ \varepsilon^{-2 \frac{k+1}{k+1-m}} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2 + \frac{\gamma-\beta}{\alpha}) \frac{k+1}{k+1-m}}, & \beta < \gamma. \end{cases}$$

- Result applies to the approximation of CDF, quantiles and CVaR with $m = 1$, and with $m = 2$ for the PDF.

Theorem (cont.)

If $\Phi \in C^{k+1}(\Theta)$ and $m \leq k$, then there exists $\hat{\Phi}_L$ such that $\text{MSE}(\hat{\Phi}_L^{(m)}) = \mathcal{O}(\varepsilon^2)$:

$$\text{comp. cost (no interp. cost)} \lesssim |\ln(\varepsilon)| \begin{cases} \varepsilon^{-2 \frac{k+1}{k+1-m}}, & \beta > \gamma, \\ \varepsilon^{-2 \frac{k+1}{k+1-m}} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2 + \frac{\gamma-\beta}{\alpha}) \frac{k+1}{k+1-m}}, & \beta < \gamma. \end{cases}$$

- Result applies to the approximation of **CDF, quantiles and CVaR** with $m = 1$, and with $m = 2$ for the PDF.
- Again, a similar complexity result is available for Φ analytic:

$$\text{comp. cost} \lesssim \varepsilon^{-2} |\ln(\varepsilon)|^{4(1+m)} + |\ln(\varepsilon)|^{3+4m} \begin{cases} \varepsilon^{-2}, & \beta > \gamma, \\ \varepsilon^{-2} \ln(\varepsilon)^2, & \beta = \gamma, \\ \varepsilon^{-(2 + \frac{\gamma-\beta}{\alpha})} |\ln(\varepsilon)|^{\frac{\gamma-\beta}{\alpha}} (1 + 2m), & \beta < \gamma. \end{cases}$$

MLMC for CDF, quantile, and CVaR: error control

- Recall that key quantities such as the CDF, quantile, and CVaR are all derived from the function

$$\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q)) \quad \text{for} \quad \phi(\vartheta, Q) = \vartheta + \frac{1}{1-\tau}(Q - \vartheta)^+.$$

- However, complexity result with $m = 1$ applies directly only to Φ' . In fact, it “only” guarantees that $\text{MSE}(\hat{\Phi}') = \mathcal{O}(\varepsilon^2)$. What about these other derived quantities?

MLMC for CDF, quantile, and CVaR: error control

- Recall that key quantities such as the CDF, quantile, and CVaR are all derived from the function

$$\Phi(\vartheta) = \mathbb{E}(\phi(\vartheta, Q)) \quad \text{for} \quad \phi(\vartheta, Q) = \vartheta + \frac{1}{1-\tau}(Q - \vartheta)^+.$$

- However, complexity result with $m = 1$ applies directly only to Φ' . In fact, it “only” guarantees that $\text{MSE}(\hat{\Phi}') = \mathcal{O}(\varepsilon^2)$. What about these other derived quantities?

Corollary

For $\tau \in (0, 1)$, let $\hat{\Phi}$ be the MLMC estimator of $\Phi \in \mathcal{C}^{k+1}(\Theta)$, $k \geq 1 = m$, so that $\text{MSE}(\hat{\Phi}') = \mathcal{O}(\varepsilon^2)$. If the true τ -quantile is an interior point of Θ , then

$$\max\{\text{quantile MSE, CVaR MSE, unif. CDF MSE}\} = \mathcal{O}(\varepsilon^2),$$

at a cost dominated by $\text{cost}(\hat{\Phi}') = \text{cost}(\text{CDF est.})$.

Corollary provides an all-at-once approach for the simultaneous approximation of CDF, quantiles, and CVaR.

Toy example: the characteristic function

- Let's consider the *toy model* to describe a **European call option** again, i.e., asset follows

$$dS = rS dt + \sigma S dW, \quad S(0) = S_0,$$

- **Quantity of interest** is the discounted “payoff”: $Q := e^{-rT} \max(S(T) - K, 0)$

Toy example: the characteristic function

- Let's consider the *toy model* to describe a **European call option** again, i.e., asset follows

$$dS = rS dt + \sigma S dW, \quad S(0) = S_0,$$

- **Quantity of interest** is the discounted “payoff”: $Q := e^{-rT} \max(S(T) - K, 0)$
- **But:** Q has a *mixed distribution*, in the sense that $\mathbb{P}(Q = 0) > 0$.
- Consequently, its CDF $F_Q := \mathbb{E}(I(Q \leq \cdot))$ has a jump discontinuity at the origin:
 \leadsto we **cannot** guarantee a uniformly accurate MLMC CDF-approximation, if $0 \in \Theta$.

Toy example: the characteristic function

- Let's consider the *toy model* to describe a **European call option** again, i.e., asset follows



$$dS = rS dt + \sigma S dW, \quad S(0) = S_0,$$

- Quantity of interest** is the discounted “payoff”: $Q := e^{-rT} \max(S(T) - K, 0)$
- But:** Q has a *mixed distribution*, in the sense that $\mathbb{P}(Q = 0) > 0$.
- Consequently, its CDF $F_Q := \mathbb{E}(I(Q \leq \cdot))$ has a jump discontinuity at the origin:
 \leadsto we **cannot** guarantee a uniformly accurate MLMC CDF-approximation, if $0 \in \Theta$.
- But we **can** guarantee an accurate MLMC approximation of the **characteristic function** to characterize probability distribution of Q :

$$\varphi_Q(\vartheta) = \underbrace{\mathbb{E}(\cos(\vartheta Q))}_{=: \phi_1(\vartheta, Q)} + i \underbrace{\mathbb{E}(\sin(\vartheta Q))}_{=: \phi_2(\vartheta, Q)} \equiv \Phi_1(\vartheta) + i \Phi_2(\vartheta),$$

- NB:** functions ϕ_i are smooth, no derivatives required (i.e., $m = 0$), moment approximations via post-processing possible [K., Nobile. 2018].

- Milstein scheme with $h_\ell = 2^{-\ell} T$; $\Theta = [-1, 1]$, $r = \frac{1}{20}$, $\sigma = \frac{1}{5}$, $T = 1$, $K = 10 = S_0$.



PDE toy example

Consider a simple Poisson equation

$$-\Delta u = f, \quad \text{in } D = (0, 1)^2,$$

with homogeneous Dirichlet boundary conditions and random forcing term f given by $f(x) = -72\xi(x_1^2 + x_2^2 - x_1 - x_2)$, $\xi \sim \chi_1^2$. **Quantity of interest:** $Q := \int_D u \, dx = \xi$.



PDE toy example

Consider a simple Poisson equation

$$-\Delta u = f, \quad \text{in } D = (0, 1)^2,$$

with homogeneous Dirichlet boundary conditions and **random forcing term** f given by $f(x) = -72\xi(x_1^2 + x_2^2 - x_1 - x_2)$, $\xi \sim \chi_1^2$. **Quantity of interest:** $Q := \int_D u \, dx = \xi$.

We approximate $\Phi(\vartheta) = \vartheta + \frac{1}{1-\tau} \mathbb{E}[(Q - \vartheta)^+]$ with $\tau = 0.95$ on $\vartheta \in \Theta = [0, 10]$ for $k = 3 \rightsquigarrow$ **theory predicts** **cost** $= \mathcal{O}(\varepsilon^{-2.3} \ln(\varepsilon^{-1}))$, appears to be conservative.



Practical bottleneck: a-priori bounds

Asymptotic complexity analysis is based on **a-priori upper bounds** for error estimates:

$$\begin{aligned} \frac{\text{MSE}(\hat{\phi}_L^{(m)})}{3} &\leq \|\phi^{(m)} - \mathcal{I}_n^{(m)}(\phi)\|_{L^\infty}^2 + \|\mathcal{I}_n^{(m)}(\phi - \phi_L)\|_{L^\infty}^2 + \mathbb{E}(\|\mathcal{I}_n^{(m)}(\phi_L - \phi_L^{\text{MLMC}})\|_{L^\infty}^2) \\ &\lesssim C_1(m)^2 n^{-2(k+1-m)} + C_2(m)^2 (n-1)^{2m} b_L^2 + C_2(m)^2 (n-1)^{2m} \log(n) \sum_{\ell=0}^L \frac{V_\ell}{N_\ell} \end{aligned}$$

where $b_L = \|\phi(\theta) - \phi_L(\theta)\|_{L^\infty}$ and $V_\ell = \text{Var}(\phi(\theta, Q_\ell) - \phi(\theta, Q_{\ell-1}))$.

Practical bottleneck: a-priori bounds


Asymptotic complexity analysis is based on **a-priori upper bounds** for error estimates:

$$\frac{\text{MSE}(\hat{\phi}_L^{(m)})}{3} \leq \|\phi^{(m)} - \mathcal{I}_n^{(m)}(\phi)\|_{L^\infty}^2 + \|\mathcal{I}_n^{(m)}(\phi - \phi_L)\|_{L^\infty}^2 + \mathbb{E}(\|\mathcal{I}_n^{(m)}(\phi_L - \phi_L^{\text{MLMC}})\|_{L^\infty}^2)$$


$$\lesssim C_1(m)^2 n^{-2(k+1-m)} + C_2(m)^2 (n-1)^{2m} b_L^2 + C_2(m)^2 (n-1)^{2m} \log(n) \sum_{\ell=0}^L \frac{V_\ell}{N_\ell}$$

where $b_L = \|\phi(\theta) - \phi_L(\theta)\|_{\ell^\infty}$ and $V_\ell = \text{Var}(\phi(\theta, Q_\ell) - \phi(\theta, Q_{\ell-1}))$.


In particular, the **inverse inequality** makes error bound **severely conservative**. For example, the **bias decay**:



$m = 0$



$m = 1$



$m = 2$

Refined a-posterior error estimators

Starting point: derive error estimators based on first error splitting directly:

- *Simplifications:* interpolation in cubic Splines

Refined a-posterior error estimators

Starting point: derive error estimators based on first error splitting directly:

- *Simplifications:* interpolation in cubic Splines
- **Error estimator for bias via bivariate kernel smoothing** of empirical measure of (Q_L, Q_{L-1}) :

$$\|\mathcal{I}_n^{(m)}(\Phi - \Phi_L)\|_{L^\infty}^2 \approx \frac{1}{s^\alpha - 1} \|\mathcal{I}_n^{(m)}(\mathbb{E}(\phi_L - \phi_{L-1}))\|_{L^\infty}$$

$$\|\mathcal{I}_n^{(m)}(\mathbb{E}(\phi_L - \phi_{L-1}))\|_{L^\infty} \approx \|\mathcal{I}_n^{(m)}(\mathbb{E}^{\text{kde}}(\phi_L - \phi_{L-1}))\|_{L^\infty}$$

where the *bivariate KDE* approximates the joint PDF of $(Q_\ell, Q_{\ell-1})$ using the N_ℓ correlated samples.

Refined a-posterior error estimators

Starting point: derive error estimators based on first error splitting directly:

- *Simplifications:* interpolation in cubic Splines
- **Error estimator for bias via bivariate kernel smoothing** of empirical measure of (Q_L, Q_{L-1}) :

$$\begin{aligned} \|\mathcal{I}_n^{(m)}(\Phi - \Phi_L)\|_{L^\infty}^2 &\approx \frac{1}{s^\alpha - 1} \|\mathcal{I}_n^{(m)}(\mathbb{E}(\phi_L - \phi_{L-1}))\|_{L^\infty} \\ \|\mathcal{I}_n^{(m)}(\mathbb{E}(\phi_L - \phi_{L-1}))\|_{L^\infty} &\approx \|\mathcal{I}_n^{(m)}(\mathbb{E}^{\text{kde}}(\phi_L - \phi_{L-1}))\|_{L^\infty} \end{aligned}$$

where the *bivariate KDE* approximates the joint PDF of $(Q_\ell, Q_{\ell-1})$ using the N_ℓ correlated samples.

- **Error estimator for statistical error via bootstrapping MLMC estimators:**

- “Observation”: an MLMC estimator is defined through the hierarchy of samples


$$\left\{ \left\{ Q_\ell^{(i,\ell)}, Q_{\ell-1}^{(i,\ell)} \right\}_{i=1}^{N_\ell} \right\}_{\ell=0}^L$$

- Idea: **resample** $N_{\text{bs}} \gg 1$ “new” MLMC estimators Ψ_j


$$\mathbb{E}(\|\mathcal{I}_n^{(m)}(\Phi_L - \Phi_L^{\text{MLMC}})\|_{L^\infty}^2) \approx \frac{1}{N_{\text{bs}}} \sum_{j=1}^{N_{\text{bs}}} \|\mathcal{I}_n^{(m)}(\Psi_j(\theta) - \bar{\Psi}(\theta))\|_{L^\infty}^2$$

Effects of refined a-posteriori error estimators


Bias estimation:



$m = 0$




$m = 1$




$m = 2$


Estimation of statistical error:



$m = 0$



$m = 1$



$m = 2$


Computational Example

In [Ayoul-Guilmarda, Ganesh, K., Nobile. 2023] we combine these with an adaptive, *continuation MLMC framework* for an efficient implementation (Python library *XMC*) that can be tuned to target CDF, VaR, or CVaR.

Computational Example

In [Ayoul-Guilmarda, Ganesh, K., Nobile. 2023] we combine these with an adaptive, *continuation MLMC framework* for an efficient implementation (Python library *xMC*) that can be tuned to target CDF, VaR, or CVaR.


Example: (steady incompressible) Navier-Stokes Flow over a Cylinder in a Channel



Computational Example

In [Ayoul-Guilmarda, Ganesh, K., Nobile. 2023] we combine these with an adaptive, *continuation MLMC framework* for an efficient implementation (Python library *XMC*) that can be tuned to target CDF, VaR, or CVaR.

Example: (steady incompressible) **Navier-Stokes Flow over a Cylinder in a Channel**



Take home message:

- we introduced a uniformly accurate MLMC estimator for a parametric expectation and its derivatives
- the approach enables approximating the characteristic function as well as CDF, PDF, VaR, and CVaR
- refined, a-posterior error estimates are required for computationally “heavy” problems when derivatives are required.

Take home message:

- we introduced a uniformly accurate MLMC estimator for a parametric expectation and its derivatives
- the approach enables approximating the characteristic function as well as CDF, PDF, VaR, and CVaR
- refined, a-posterior error estimates are required for computationally “heavy” problems when derivatives are required.

Thank you for your attention.

Further details:

SK and F. Nobile. Multilevel Monte Carlo Approximation of Functions. *SIAM/ASA J. Uncertain. Quantif.*, **3**(6):1256–1293, 2018.

Q. Ayoul-Guilmard, S. Ganesh, SK, and F. Nobile. Quantifying uncertain system outputs via the multi-level Monte Carlo method – distribution and robustness measures. *Int. J. Uncertain. Quantif.*, **13**(5):61–98, 2023.

References



Q. Ayoul-Guilmard, S. Ganesh, S. Krumscheid, and F. Nobile.
Quantifying uncertain system outputs via the multi-level Monte Carlo method – distribution and robustness measures.
Int. J. Uncertain. Quantif., 13(5):61–98, 2023.



S. Krumscheid, F. Nobile, and M. Pisaroni.
Quantifying uncertain system outputs via the multilevel Monte Carlo method — Part I: Central moment estimation.
J. Comput. Phys., 414:109466, 2020.



S. Krumscheid and F. Nobile.
Multilevel Monte Carlo Approximation of Functions.
SIAM/ASA J. Uncertain. Quantif., 3(6):1256–1293, 2018.



H. Hoel and S. Krumscheid.
Central limit theorems for multilevel Monte Carlo methods.
J. Complexity, 54:101407, 2019.



N. Collier, A.-L. Haji-Ali, F. Nobile, E. von Schwerin, and R. Tempone
A continuation multilevel Monte Carlo algorithm
Bit Numer. Math., 55:399–432, 2015.



ExaQute consortium.
XMC Python library, <http://doi.org/10.5281/zenodo.3235833>.

References II



M.B. Giles.

Multilevel Monte Carlo methods.

Acta Numer., 24:259–328, 2015.



C. Bierig and A. Chernov.

Estimation of arbitrary order central statistical moments by the multilevel Monte Carlo method.

Stoch. PDE: Anal. Comp., 4(1):3–40, 2016.



C. Bierig and A. Chernov.

Convergence analysis of multilevel Monte Carlo variance estimators and application for random obstacle problems.

Numer. Math., 130(4):579–613, 2015.