

Uniform in time approximations

M. Ottobre (Heriot Watt University and Maxwell Institute)

Joint work with D. Crisan, E. Zatorska (Imperial), P. Dobson (Delft),
B. Goddard (Edinburgh), L. Angeli, I. Souttar (Heriot Watt), Julien
Barre' (Orleans)

MCMC2023

Overview

- ▶ What I mean by ‘uniform in time approximation’

Overview

- ▶ What I mean by ‘uniform in time approximation’
- ▶ General approach

Overview

- ▶ What I mean by ‘uniform in time approximation’
- ▶ General approach
- ▶ Several instances to which the approach is applicable

Overview

- ▶ What I mean by ‘uniform in time approximation’
- ▶ General approach
- ▶ Several instances to which the approach is applicable
- ▶ Strengths and limitations

Context: UiT approximations

- Suppose X_t solves a (one-dimensional, elliptic) SDE

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x,$$

Context: UiT approximations

- ▶ Suppose X_t solves a (one-dimensional, elliptic) SDE

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x,$$

- ▶ Let Y_t be "some approximation", for example numerical approximation Y_t^δ

Context: UiT approximations

- ▶ Suppose X_t solves a (one-dimensional, elliptic) SDE

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x,$$

- ▶ Let Y_t be "some approximation", for example numerical approximation Y_t^δ
- ▶ We want sufficient criteria to guarantee

$$|\mathbb{E}f(X_t) - \mathbb{E}f(Y_t^\delta)| \leq K\delta, \quad \forall f \in \mathcal{C}_b(\mathbb{R})$$

with K independent of time (but dependent on f).

Context: UiT approximations

- ▶ Suppose X_t solves a (one-dimensional, elliptic) SDE

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x,$$

- ▶ Let Y_t be "some approximation", for example numerical approximation Y_t^δ
- ▶ We want sufficient criteria to guarantee

$$|\mathbb{E}f(X_t) - \mathbb{E}f(Y_t^\delta)| \leq K\delta, \quad \forall f \in \mathcal{C}_b(\mathbb{R})$$

with K independent of time (but dependent on f).

Some words of warning

Some words of warning

- ▶ **This is not always possible!** Whether it is possible or not depends on

Some words of warning

- ▶ **This is not always possible!** Whether it is possible or not depends on
 1. The dynamics to approximate
 2. The approximation

Some words of warning

- ▶ **This is not always possible!** Whether it is possible or not depends on
 1. The dynamics to approximate
 2. The approximation
- ▶ I am not considering time-averages (though you get them as a consequence - modulo having an invariant measure)

Some words of warning

- ▶ **This is not always possible!** Whether it is possible or not depends on
 1. The dynamics to approximate
 2. The approximation
- ▶ I am not considering time-averages (though you get them as a consequence - modulo having an invariant measure)
- ▶ I don't necessarily want to approximate the invariant measure, I want to approximate the law at time t , for every $t > 0$

How do we go about this?

Setting

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x.$$

Setting

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x.$$

► $\mathbb{E}f(X_t^x) =: u(t, x)$

Setting

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x.$$

► $\mathbb{E}f(X_t^x) =: u(t, x)$

$$u(t, x) = (\mathcal{P}_t f)(x) \quad \text{Markov semigroup}$$

Setting

$$dX_t = b(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x.$$

► $\mathbb{E}f(X_t^x) =: u(t, x)$

$$u(t, x) = (\mathcal{P}_t f)(x) \quad \text{Markov semigroup}$$

► $u(t, x)$ solves a PDE

$$\partial_t u(t, x) = \mathcal{L}u(t, x) \quad u(0, x) = f(x).$$

$$\mathcal{L} = b(x)\partial_x + \frac{1}{2}\sigma^2(x)\partial_{xx}$$

General Approach

General Approach

$$\begin{aligned} & \mathbb{E}f(X_t) - \mathbb{E}f(Y_t^\delta) \\ = & (\mathcal{P}_t f)(x) - (\mathcal{P}_t^\delta f)(x) \\ = & \int_0^t ds \mathcal{P}_{t-s}^\delta (\mathcal{L} - \mathcal{L}^\delta) \mathcal{P}_s f(x, y) \\ \leq & \int_0^t \|(\mathcal{L} - \mathcal{L}^\delta)(\mathcal{P}_s f)\|_\infty ds \\ \simeq & \delta \int_0^t \|\partial_x^2 (\mathcal{P}_s f)\|_\infty ds \end{aligned}$$

Key stability condition

$$\boxed{|\partial_x(\mathcal{P}_t f)(x)| \leq c e^{-\lambda t}} \quad \text{for some } \lambda > 0 \quad (\text{DE})$$

Key stability condition

$$\boxed{|\partial_x(\mathcal{P}_t f)(x)| \leq c e^{-\lambda t}} \quad \text{for some } \lambda > 0 \quad (\text{DE})$$

► if the above is true then

$$|\mathbb{E}f(X_t) - \mathbb{E}f(Y_t^\delta)| \leq \delta \int_0^t \|\partial_x^2(\mathcal{P}_s f)\|_\infty ds \leq \delta K \int_0^t e^{-\lambda s} ds \leq \delta K.$$

Key stability condition

$$\boxed{|\partial_x(\mathcal{P}_t f)(x)| \leq c e^{-\lambda t}} \quad \text{for some } \lambda > 0 \quad (\text{DE})$$

- ▶ if the above is true then

$$|\mathbb{E}f(X_t) - \mathbb{E}f(Y_t^\delta)| \leq \delta \int_0^t \|\partial_x^2(\mathcal{P}_s f)\|_\infty ds \leq \delta K \int_0^t e^{-\lambda s} ds \leq \delta K.$$

- ▶ Why is this a stability condition?

Key stability condition

$$\boxed{|\partial_x(\mathcal{P}_t f)(x)| \leq c e^{-\lambda t}} \quad \text{for some } \lambda > 0 \quad (\text{DE})$$

- ▶ if the above is true then

$$|\mathbb{E}f(X_t) - \mathbb{E}f(Y_t^\delta)| \leq \delta \int_0^t \|\partial_x^2(\mathcal{P}_s f)\|_\infty ds \leq \delta K \int_0^t e^{-\lambda s} ds \leq \delta K.$$

- ▶ Why is this a stability condition?

$$\int_0^\ell \frac{d}{du}(\mathcal{P}_t f)(\gamma(u)) du = (\mathcal{P}_t f)(y) - (\mathcal{P}_t f)(x) = \int_0^\ell (\partial_x \mathcal{P}_t)(\gamma(u)) du$$

Averaging

- Fast-slow dynamics, $(X_t, Y_t) \in \mathbb{R}^d \times \mathbb{R}^n$

$$dX_t = f(X_t, Y_t)dt + g(X_t, Y_t)dW_t$$

$$dY_t = \frac{1}{\epsilon}h(X_t, Y_t)dt + \frac{1}{\sqrt{\epsilon}}\sigma(X_t, Y_t)dB_t, \quad 0 < \epsilon \ll 1.$$

Averaging

- Fast-slow dynamics, $(X_t, Y_t) \in \mathbb{R}^d \times \mathbb{R}^n$

$$dX_t = f(X_t, Y_t)dt + g(X_t, Y_t)dW_t$$

$$dY_t = \frac{1}{\epsilon}h(X_t, Y_t)dt + \frac{1}{\sqrt{\epsilon}}\sigma(X_t, Y_t)dB_t, \quad 0 < \epsilon \ll 1.$$

- Limiting dynamics

$$d\bar{X}_t = F(\bar{X}_t)dt + G(\bar{X}_t)dW_t$$

where

$$F(x) = \int f(x, y)\mu^x(dy), \quad G(x) = \int g(x, y)\mu^x(dy)$$

Averaging

- Fast-slow dynamics, $(X_t, Y_t) \in \mathbb{R}^d \times \mathbb{R}^n$

$$dX_t = f(X_t, Y_t)dt + g(X_t, Y_t)dW_t$$

$$dY_t = \frac{1}{\epsilon}h(X_t, Y_t)dt + \frac{1}{\sqrt{\epsilon}}\sigma(X_t, Y_t)dB_t, \quad 0 < \epsilon \ll 1.$$

- Limiting dynamics

$$d\bar{X}_t = F(\bar{X}_t)dt + G(\bar{X}_t)dW_t$$

where

$$F(x) = \int f(x, y)\mu^x(dy), \quad G(x) = \int g(x, y)\mu^x(dy)$$

- Conceptual problems in practice if you don't have UiT approximation

- ▶ Associated Poisson equation:

$$(L^x f)(x, y) = \varphi(x, y)$$

- ▶ L^x is the generator of the fast process

$$dY_t = h(X_t, Y_t)dt + \sigma(X_t, Y_t)dB_t$$

namely

$$L^x f = h(x, y)\partial_y f + \sigma^2(x, y)\partial_{yy} f$$

$$L^x f = \varphi$$

- ▶ Representation formula

$$f(x, y) = \int_0^\infty (P_t^x \varphi)(y) dt$$

- ▶ smoothness of solution, in both x and y
- ▶ quantify how solution $f(x, y)$ varies as x varies - similarly for μ^x

Multiscale Problems for Interacting Particle Systems

- ▶ Particle system interacting with fast network

$$dX_t^i = -\nabla V(X_t^i)dt + \sum_{j \neq i} A_{ij}^\varepsilon(t) K(X_t^i - X_t^j)dt + \sqrt{2D}dB_t^i, \quad i = 1 \dots N$$

$$dA_{ij}^\varepsilon(t) = -A_{ij}^\varepsilon(t_-)dN_{ij}^{d,\varepsilon}(t) + [1 - A_{ij}^\varepsilon(t_-)]\mathbf{1}_{\{|X_t^i - X_t^j| \leq R\}}dN_{ij}^{f,\varepsilon}(t)$$

Multiscale Problems for Interacting Particle Systems

- ▶ Particle system interacting with fast network

$$dX_t^i = -\nabla V(X_t^i)dt + \sum_{j \neq i} A_{ij}^\varepsilon(t) K(X_t^i - X_t^j)dt + \sqrt{2D}dB_t^i, \quad i = 1 \dots N$$

$$dA_{ij}^\varepsilon(t) = -A_{ij}^\varepsilon(t_-)dN_{ij}^{d,\varepsilon}(t) + [1 - A_{ij}^\varepsilon(t_-)]\mathbf{1}_{\{|X_t^i - X_t^j| \leq R\}}dN_{ij}^{f,\varepsilon}(t)$$

- ▶ Main features:
 - ▶ large number of particles $N \rightarrow \infty$ and fast network evolution $\varepsilon \rightarrow 0$
 - ▶ the network is *sparse*, i.e. the interaction is not of mean-field type

Are Derivative estimates hard to obtain?

Are Derivative estimates hard to obtain?

By now various simple criteria

- Numerical methods

- [1] D. Crisan, P. Dobson, M. O. *Uniform in time estimates for the weak error of the Euler method*, Trans. AMS.
- [2] L. Angeli, D. Crisan, M.O. *Uniform in time convergence of numerical schemes for SDEs via Strong Exponential Stability: Euler methods, Split-Step and Tamed Schemes*. arxiv

- Particle systems

- [3] J. Barre, P. Dobson, M. O., E. Zatorska. *Non mean field fast network: uniform in time averaging*, SIAM Analysis

- Averaging

- [4] B. Goddard, M. Ottobre, P. Dobson, I. Souttar *Poisson equations and uniform in time averaging* arxiv

- Analytic approach

- ▶ S. Mischler, C. Mouhot. *Kac's programme in kinetic theory*, Inventiones, 2013

- Probabilistic approach

- ▶ S. Mischler, C. Mouhot. *Kac's programme in kinetic theory*, Inventiones, 2013
- ▶ D. Heydecker. *Convergence of the hard spheres Kac Process*. AOP, 2019
- ▶ A. Durmus, A. Eberle
- ▶ Series of papers by P. Dobson, J. Sanz-Serna, K. Zygalakis