

Infinite-dimensional Wishart processes

Sonja Cox, Christa Cuchiero, and Asma Khedher

MCM 2023, Paris



UNIVERSITY OF AMSTERDAM

The Wishart distribution

Let

- ▶ $n, p \in \mathbb{N}$,
- ▶ $\Gamma = (\gamma_{i,j})_{1 \leq i \leq p, 1 \leq j \leq n}$ be i.i.d. standard Gaussian random variables,
- ▶ $S^+(\mathbb{R}^p) = \{A \in \mathbb{R}^{p \times p} : A \text{ is positive semi-definite}\}$,
- ▶ $S^{++}(\mathbb{R}^p) = \{A \in \mathbb{R}^{p \times p} : A \text{ is positive definite}\}$.

¹The Wishart distribution was introduced in 1928 by John Wishart

The Wishart distribution

Let

- ▶ $n, p \in \mathbb{N}$,
- ▶ $\Gamma = (\gamma_{i,j})_{1 \leq i \leq p, 1 \leq j \leq n}$ be i.i.d. standard Gaussian random variables,
- ▶ $S^+(\mathbb{R}^p) = \{A \in \mathbb{R}^{p \times p} : A \text{ is positive semi-definite}\}$,
- ▶ $S^{++}(\mathbb{R}^p) = \{A \in \mathbb{R}^{p \times p} : A \text{ is positive definite}\}$.

Let X be the $S^+(\mathbb{R}^p)$ -valued random variable given by

$$X = \Gamma \Gamma^T,$$

then¹ $X \sim \text{Wishart}_p(n)$ (' X had a Wishart distribution with n degrees of freedom').

¹The Wishart distribution was introduced in 1928 by John Wishart

The Wishart distribution

Let

- ▶ $n, p \in \mathbb{N}$,
- ▶ $\Gamma = (\gamma_{i,j})_{1 \leq i \leq p, 1 \leq j \leq n}$ be i.i.d. standard Gaussian random variables,
- ▶ $S^+(\mathbb{R}^p) = \{A \in \mathbb{R}^{p \times p} : A \text{ is positive semi-definite}\}$,
- ▶ $S^{++}(\mathbb{R}^p) = \{A \in \mathbb{R}^{p \times p} : A \text{ is positive definite}\}$.

Let X be the $S^+(\mathbb{R}^p)$ -valued random variable given by

$$X = \Gamma \Gamma^T,$$

then¹ $X \sim \text{Wishart}_p(n)$ (' X had a Wishart distribution with n degrees of freedom'). Generalizations:

$$X = (\sqrt{Q}\Gamma + A)(\sqrt{Q}\Gamma + A)^T$$

with $Q \in S^+(\mathbb{R}^p)$, $A \in \mathbb{R}^{p \times n}$: $X \sim \text{Wishart}_p(n, Q, A)$.

¹The Wishart distribution was introduced in 1928 by John Wishart

The Wishart distribution

- Let $X \sim \text{Wishart}_p(n)$. If $n \geq p$ then $X \in S^{++}(\mathbb{R}^p)$ a.s. and X allows for a density on $S^{++}(\mathbb{R}^p)$.

The Wishart distribution

- ▶ Let $X \sim \text{Wishart}_p(n)$. If $n \geq p$ then $X \in S^{++}(\mathbb{R}^p)$ a.s. and X allows for a density on $S^{++}(\mathbb{R}^p)$.
- ▶ The Laplace transform $S^+(\mathbb{R}^p) \ni u \mapsto \mathbb{E}(e^{-\text{tr}(uX)})$ can be given explicitly.

The Wishart distribution

- ▶ Let $X \sim \text{Wishart}_p(n)$. If $n \geq p$ then $X \in S^{++}(\mathbb{R}^p)$ a.s. and X allows for a density on $S^{++}(\mathbb{R}^p)$.
- ▶ The Laplace transform $S^+(\mathbb{R}^p) \ni u \mapsto \mathbb{E}(e^{-\text{tr}(uX)})$ can be given explicitly.
- ▶ The Wishart distribution is used in e.g. multidimensional (Bayesian) statistical analysis and random matrix theory.

The Wishart process

Let

- ▶ $A \in \mathbb{R}^{p \times p}$,
- ▶ $Y_0, Q \in S^{++}(\mathbb{R}^p)$,
- ▶ $W: [0, \infty) \times \Omega \rightarrow \mathbb{R}^{p \times n}$ a standard Brownian motion,

and let $Y: [0, \infty) \times \Omega \rightarrow \mathbb{R}^{p \times n}$ be the Ornstein-Uhlenbeck process satisfying

$$dY_t = AY_t dt + \sqrt{Q} dW_t, \quad t \geq 0,$$

$$\text{i.e., } Y_t = e^{tA}Y_0 + \int_0^t e^{(t-s)A}\sqrt{Q} dW_s.$$

The Wishart process

Let

- ▶ $A \in \mathbb{R}^{p \times p}$,
- ▶ $Y_0, Q \in S^{++}(\mathbb{R}^p)$,
- ▶ $W: [0, \infty) \times \Omega \rightarrow \mathbb{R}^{p \times n}$ a standard Brownian motion,

and let $Y: [0, \infty) \times \Omega \rightarrow \mathbb{R}^{p \times n}$ be the Ornstein-Uhlenbeck process satisfying

$$dY_t = AY_t dt + \sqrt{Q} dW_t, \quad t \geq 0,$$

i.e., $Y_t = e^{tA}Y_0 + \int_0^t e^{(t-s)A}\sqrt{Q} dW_s$. Set

$$X_t = Y_t Y_t^T, \quad t \geq 0.$$

Then X_t has a Wishart distribution for all $t > 0$. We call $(X_t)_{t \geq 0}$ a **Wishart process**.

The Wishart process

Theorem (Bru, 1989)

Let $(X_t)_{t \geq 0}$ be the Wishart process introduced above and $n \geq p$. Then there exists an $\mathbb{R}^{p \times p}$ -standard Brownian motion $(B_t)_{t \geq 0}$ such that

$$dX_t = (nQ + AX_t + X_t A^T) dt + \sqrt{Q} dB_t X_t + X_t dB_t^T \sqrt{Q}, \quad t \geq 0.$$

Proof.

Apply Itô formula to $Y_t Y_t^T$. Show that martingale part of $dY_t Y_t^T$ is equal in distribution to $\sqrt{Q} dB_t X_t + X_t dB_t^T \sqrt{Q}$. \square

The Wishart process

Theorem (Bru, 1991)

If $\alpha \in \mathbb{N} \cup (p - 1, \infty)$ and $\text{rank}(X_0) \leq \alpha$ then there exists a unique $S^+(\mathbb{R}^p)$ -valued (weak) solution to

$$dX_t = \alpha Q + AX_t + X_t A^T dt \sqrt{Q} dB_t X_t + X_t dB_t^T \sqrt{Q}, \quad t \geq 0.$$

(Wis_p)

The Wishart process

Theorem (Bru, 1991)

If $\alpha \in \mathbb{N} \cup (p - 1, \infty)$ and $\text{rank}(X_0) \leq \alpha$ then there exists a unique $S^+(\mathbb{R}^p)$ -valued (weak) solution to

$$dX_t = \alpha Q + AX_t + X_t A^T dt \sqrt{Q} dB_t X_t + X_t dB_t^T \sqrt{Q}, \quad t \geq 0. \quad (\text{Wis}_p)$$

Moreover,

- ▶ *the Laplace transform of X_t , $t \geq 0$, can be given explicitly and coincides with a Wishart distribution when $\alpha \in \mathbb{N}$;*
- ▶ *the eigenvalues and -vectors of X_t , $t \geq 0$, solve a system of SDEs that can be given explicitly.*

The Wishart process

Theorem (Bru, 1991)

If $\alpha \in \mathbb{N} \cup (p - 1, \infty)$ and $\text{rank}(X_0) \leq \alpha$ then there exists a unique $S^+(\mathbb{R}^p)$ -valued (weak) solution to

$$dX_t = \alpha Q + AX_t + X_t A^T dt + \sqrt{Q} dB_t X_t + X_t dB_t^T \sqrt{Q}, \quad t \geq 0. \quad (\text{Wis}_p)$$

Moreover,

- ▶ *the Laplace transform of X_t , $t \geq 0$, can be given explicitly and coincides with a Wishart distribution when $\alpha \in \mathbb{N}$;*
- ▶ *the eigenvalues and -vectors of X_t , $t \geq 0$, solve a system of SDEs that can be given explicitly.*

Theorem (Cuchiero, Filipovic, Mayerhofer, Teichman: 2011)

If a (weak) solution to (Wis_p) exists for all $X_0 \in S^{++}(\mathbb{R}^p)$ then $\alpha > p - 1$.

Infinite dimensional Wishart processes: why?


Infinite-dimensional stochastic processes call for infinite-dimensional covariance models.

Infinite dimensional Wishart processes: why?

Infinite-dimensional stochastic processes call for infinite-dimensional covariance models.

E.g.: models in financial/energy/commodity markets, i.e., models describing, **for every** $t_0 \leq t \leq t_1$, the expected cost of buying 1 euro/kWh/kg of silver, given the information at time t_0 .

Samples of the forward rate curve f_t for different values of t



Question: can we construct infinite-dimensional Wishart processes?

Question: can we construct infinite-dimensional Wishart processes?

Recall:

Question: can we construct infinite-dimensional Wishart processes?

Recall:

Theorem (Bru, 1991)

If $\alpha \in \mathbb{N} \cup (p - 1, \infty)$ and $\text{rank}(X_0) \leq \alpha$ then there exists a unique $S^+(\mathbb{R}^p)$ -valued (stoch. weak) solution to

$$dX_t = \alpha Q + AX_t + X_t A^T dt \sqrt{Q} dB_t X_t + X_t dB_t^T \sqrt{Q}, \quad t \geq 0. \\ (\text{Wis}_p)$$

Theorem (Cuchiero, Filipovic, Mayerhofer, Teichman: 2011)

If a (weak) solution to (Wis_p) exists for all $X_0 \in S^{++}(\mathbb{R}^p)$ then $\alpha > p - 1$.

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,
- ▶ $L_1(H)$ the space of trace-class operators on H ,

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,
- ▶ $L_1(H)$ the space of trace-class operators on H ,
- ▶ $L_2(H)$ the space of Hilbert-Schmidt operators on H ,

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,
- ▶ $L_1(H)$ the space of trace-class operators on H ,
- ▶ $L_2(H)$ the space of Hilbert-Schmidt operators on H ,
- ▶ $S(H) = \{A \in L(H) : A^* = A\}$,

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,
- ▶ $L_1(H)$ the space of trace-class operators on H ,
- ▶ $L_2(H)$ the space of Hilbert-Schmidt operators on H ,
- ▶ $S(H) = \{A \in L(H): A^* = A\}$,
- ▶ $S^+(H) = \{A \in S(H): A \geq 0\}$,

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,
- ▶ $L_1(H)$ the space of trace-class operators on H ,
- ▶ $L_2(H)$ the space of Hilbert-Schmidt operators on H ,
- ▶ $S(H) = \{A \in L(H) : A^* = A\}$,
- ▶ $S^+(H) = \{A \in S(H) : A \geq 0\}$,
- ▶ $S_1(H) = S(H) \cap L_1(H)$,

Some notation

- ▶ $(H, \langle \cdot, \cdot \rangle_H, \|\cdot\|_H)$ a separable real Hilbert space,
- ▶ $L_1(H)$ the space of trace-class operators on H ,
- ▶ $L_2(H)$ the space of Hilbert-Schmidt operators on H ,
- ▶ $S(H) = \{A \in L(H) : A^* = A\}$,
- ▶ $S^+(H) = \{A \in S(H) : A \geq 0\}$,
- ▶ $S_1(H) = S(H) \cap L_1(H)$,
- ▶ $S_1^+(H) = S^+(H) \cap L_1(H)$.

Infinite-dimensional Wishart processes: an example

Benth and Simonsen (2018) consider an 'infinite-dimensional Heston model':

Infinite-dimensional Wishart processes: an example

Benth and Simonsen (2018) consider an 'infinite-dimensional Heston model':

the stochastic covariance is modelled by $X_t := Y_t \otimes Y_t$, where Y is an H -valued Ornstein-Uhlenbeck process:

$$dY_t = AY_t dt + \sqrt{Q}dW_t,$$

with

- ▶ $A: D(A) \subseteq H \rightarrow H$ is the generator of a C_0 -semigroup,
- ▶ W is a H -cylindrical Brownian motion,
- ▶ $Q \in S_1^+(H)$.

Infinite-dimensional Wishart processes: an example

Benth and Simonsen (2018) consider an ‘infinite-dimensional Heston model’:

the stochastic covariance is modelled by $X_t := Y_t \otimes Y_t$, where Y is an H -valued Ornstein-Uhlenbeck process:

$$dY_t = AY_t dt + \sqrt{Q}dW_t,$$

with

- ▶ $A: D(A) \subseteq H \rightarrow H$ is the generator of a C_0 -semigroup,
- ▶ W is a H -cylindrical Brownian motion,
- ▶ $Q \in S_1^+(H)$.

Note: $X_t \in S_1^+(H)$ a.s.

Infinite-dimensional Wishart processes: an example

Benth and Simonsen (2018) consider an ‘infinite-dimensional Heston model’:

the stochastic covariance is modelled by $X_t := Y_t \otimes Y_t$, where Y is an H -valued Ornstein-Uhlenbeck process:

$$dY_t = AY_t dt + \sqrt{Q}dW_t,$$

with

- ▶ $A: D(A) \subseteq H \rightarrow H$ is the generator of a C_0 -semigroup,
- ▶ W is a H -cylindrical Brownian motion,
- ▶ $Q \in S_1^+(H)$.

Note: $X_t \in S_1^+(H)$ a.s.

Indeed, Benth and Simonsen calculate the Laplace transform

$$S_1^+(H) \ni u \mapsto \mathbb{E}e^{-\langle uY_t, Y_t \rangle_H}.$$

This Laplace transform is analogous to that of a Wishart distribution with 1 degree of freedom.

Theorem (C., Cuchiero, Khedher (2023))

Assume

- ▶ $A: D(A) \subseteq H \rightarrow H$ generator of a C_0 -semigroup $(e^{tA})_{t \geq 0}$,
- ▶ $Q \in S^+(H)$ such that $\int_0^t \|Qe^{sA}\|_{L_2(H)} ds < \infty$ for all $t > 0$,
- ▶ $n \in \mathbb{N}$ and $X_0 \in S(H)$ with $\text{rank}(X_0) \leq n$.

Then there exists a $L_2(H)$ -cylindrical Brownian motion W and an adapted stochastic process $X: [0, \infty) \times \Omega \rightarrow S_1^+(H)$ such that $\forall g, h \in D(A^), \forall t > 0$:*

$$\begin{aligned} d\langle X_t g, h \rangle_H = & (n\langle Qg, h \rangle_H + \langle A^* g, X_t h \rangle_H + \langle X_t g, A^* h \rangle_H) dt \\ & + \langle \sqrt{X_t} dW_t \sqrt{Q} g, h \rangle_H + \langle \sqrt{Q} dW_t^* \sqrt{X_t} g, h \rangle_H. \end{aligned}$$

(Wis $_{\infty}$)

Theorem (C., Cuchiero, Khedher (2023))

Assume

- ▶ $A: D(A) \subseteq H \rightarrow H$ generator of a C_0 -semigroup $(e^{tA})_{t \geq 0}$,
- ▶ $Q \in S^+(H)$ such that $\int_0^t \|Qe^{sA}\|_{L_2(H)} ds < \infty$ for all $t > 0$,
- ▶ $n \in \mathbb{N}$ and $X_0 \in S(H)$ with $\text{rank}(X_0) \leq n$.

Then there exists a $L_2(H)$ -cylindrical Brownian motion W and an adapted stochastic process $X: [0, \infty) \times \Omega \rightarrow S_1^+(H)$ such that $\forall g, h \in D(A^), \forall t > 0$:*

$$\begin{aligned} d\langle X_t g, h \rangle_H = & (n\langle Qg, h \rangle_H + \langle A^* g, X_t h \rangle_H + \langle X_t g, A^* h \rangle_H) dt \\ & + \langle \sqrt{X_t} dW_t \sqrt{Q} g, h \rangle_H + \langle \sqrt{Q} dW_t^* \sqrt{X_t} g, h \rangle_H. \end{aligned}$$

(Wis_∞)

Moreover, this process is unique in law.

Existence.

Let $Y: [0, \infty) \times \Omega \rightarrow L(\mathbb{R}^n, H)$ satisfy $Y_0 Y_0^* = X_0$ and

$$Y_t = e^{tA} Y_0 + \int_0^t e^{(t-s)A} \sqrt{Q} d\tilde{W}_s$$

where \tilde{W} is an $L_2(\mathbb{R}^n, H)$ -cylindrical Brownian motion. Then there exists an $L_2(H)$ -cylindrical motion W such that

$$X = YY^*$$

satisfies $\forall g, h \in D(A), \forall t \geq 0$:

$$\begin{aligned} d\langle X_t g, h \rangle_H &= (n\langle Qg, h \rangle_H + \langle A^* g, X_t h \rangle_H + \langle X_t g, A^* h \rangle_H) dt \\ &\quad + \langle \sqrt{X_t} dW_t \sqrt{Q} g, h \rangle_H + \langle \sqrt{Q} dW_t^* \sqrt{X_t} g, h \rangle_H. \end{aligned}$$



Remark regarding existence

Let

$$\tilde{W}^{(i)} = \tilde{W}e_i, \quad i \in \{1, \dots, n\},$$

where e_i is the i^{th} unit vector in \mathbb{R}^n

(recall: \tilde{W} is an $L_2(\mathbb{R}_n, H)$ -cylindrical Brownian motion).

Note: $\tilde{W}^{(1)}, \dots, \tilde{W}^{(n)}$ are independent H -cylindrical Brownian motions.

Remark regarding existence

Let

$$\tilde{W}^{(i)} = \tilde{W}e_i, \quad i \in \{1, \dots, n\},$$

where e_i is the i^{th} unit vector in \mathbb{R}^n

(recall: \tilde{W} is an $L_2(\mathbb{R}_n, H)$ -cylindrical Brownian motion).

Note: $\tilde{W}^{(1)}, \dots, \tilde{W}^{(n)}$ are independent H -cylindrical Brownian motions.

Let $Y^{(1)}, \dots, Y^{(n)}: [0, \infty) \times \Omega \rightarrow H$ satisfy

$$Y_t^{(i)} = e^{tA}Y_0e_i + \int_0^t e^{(t-s)A}\sqrt{Q}dW_s^{(i)}, \quad t \geq 0.$$

Then

$$X_t = \sum_{i=1}^n Y_t^{(i)} \otimes Y_t^{(i)}, \quad t \geq 0.$$

Uniqueness in law.

Suppose $(X_t)_{t \geq 0}$ satisfies (Wis_∞) . Fix $t > 0$, $u \in S^+(H)$, $v \in S(H)$. By applying Itô's formula to

$$\Phi(s, X_s) = \exp \left(-\text{tr}(\psi(t-s, u-iv)X_s) - n \int_0^{t-s} \text{tr}(\psi(r, u-iv)Q)dr \right)$$

where $\psi: [0, t] \times L(H_{\mathbb{C}}) \rightarrow L(H_{\mathbb{C}})$ is an unknown function, we obtain that if ψ satisfies (Ricc) below then

$$\mathbb{E}(e^{-\text{tr}((u-iv)X_t)}) = \exp(-\text{tr}(\psi(t, u-iv)x_0) - \phi(t-s, u-iv))$$

Uniqueness in law.

Suppose $(X_t)_{t \geq 0}$ satisfies (Wis_∞) . Fix $t > 0$, $u \in S^+(H)$, $v \in S(H)$. By applying Itô's formula to

$$\Phi(s, X_s) = \exp \left(-\text{tr}(\psi(t-s, u-iv)X_s) - n \int_0^{t-s} \text{tr}(\psi(r, u-iv)Q)dr \right)$$

where $\psi: [0, t] \times L(H_{\mathbb{C}}) \rightarrow L(H_{\mathbb{C}})$ is an unknown function, we obtain that if ψ satisfies (Ricc) below then

$$\mathbb{E}(e^{-\text{tr}((u-iv)X_t)}) = \exp(-\text{tr}(\psi(t, u-iv)x_0) - \phi(t-s, u-iv))$$

$$\begin{cases} \frac{\partial}{\partial t} \psi(t, u-iv) = A^* \psi(t, u-iv) + \psi(t, u-iv)A \\ \quad - \frac{1}{2}(\psi(t, u-iv) + \psi^T(t, u-iv))Q(\psi(t, u-iv) + \psi^T(t, u-iv)), & t \in [0, \infty); \\ \psi(0, u-iv) = u-iv. \end{cases}$$

(Ricc)



Remarks Laplace transform

- ▶ As X is a analytically weak solution and (Ricc) typically does not allow for a strong solution some nasty approximation arguments are needed.

Remarks Laplace transform

- ▶ As X is a analytically weak solution and (Ricc) typically does not allow for a strong solution some nasty approximation arguments are needed.
- ▶ We obtain an expression for $\mathbb{E}(e^{-\text{tr}((u-iv)X_t)})$ in each of the following situations
 - ▶ $u \in S^+(H)$, $v = 0$;
 - ▶ $u = 0$, $v \in S^+(H)$ or $-v \in S^+(H)$;
 - ▶ $u \in S^+(H)$ and $v \in S(H)$ sufficiently small.

Remarks Laplace transform

- ▶ As X is an analytically weak solution and (Ricc) typically does not allow for a strong solution some nasty approximation arguments are needed.
- ▶ We obtain an expression for $\mathbb{E}(e^{-\operatorname{tr}((u-iv)X_t)})$ in each of the following situations
 - ▶ $u \in S^+(H)$, $v = 0$;
 - ▶ $u = 0$, $v \in S^+(H)$ or $-v \in S^+(H)$;
 - ▶ $u \in S^+(H)$ and $v \in S(H)$ sufficiently small.
- ▶ We can replace $n \in \mathbb{N}$ by $\alpha \in \mathbb{R}$.

Remarks Laplace transform

- ▶ As X is a analytically weak solution and (Ricc) typically does not allow for a strong solution some nasty approximation arguments are needed.
- ▶ We obtain an expression for $\mathbb{E}(e^{-\text{tr}((u-iv)X_t)})$ in each of the following situations
 - ▶ $u \in S^+(H)$, $v = 0$;
 - ▶ $u = 0$, $v \in S^+(H)$ or $-v \in S^+(H)$;
 - ▶ $u \in S^+(H)$ and $v \in S(H)$ sufficiently small.
- ▶ We can replace $n \in \mathbb{N}$ by $\alpha \in \mathbb{R}$.
- ▶ Concrete example: for $u \in S^+(H)$ we get

$$\begin{aligned} & \mathbb{E}[\exp(-\text{tr}(uX_t)) \mid x_0] \\ &= \det(I_H + 2Q_{t,u})^{-\frac{n}{2}} \exp\left(-\text{tr}\left(e^{tA^*} \sqrt{u}(I_H + 2Q_{t,u})^{-1} \sqrt{u} e^{tA} x_0\right)\right), \end{aligned}$$

$$\text{with } Q_{t,u} = \sqrt{u} \int_0^t e^{sA} Q e^{sA^*} ds \sqrt{u}.$$

Theorem (C., Cuchiero, Khedher (2023))

Assume

- ▶ $A: D(A) \subseteq H \rightarrow H$ generator of a C_0 -semigroup $(e^{tA})_{t \geq 0}$ **and** e^{tA} **is injective for some $t > 0$,**
- ▶ $Q \in S^+(H)$ **is injective** and $\int_0^t \|Qe^{sA}\|_{L_2(H)} ds < \infty$ for all $t > 0$,
- ▶ $\alpha \in \mathbb{R}$,
- ▶ W is an $L_2(H)$ -cylindrical Brownian motion W ,
- ▶ $X: [0, \infty) \times \Omega \rightarrow S_1^+(H)$ such that $\forall g, h \in D(A^*), \forall t \geq 0$:

$$\begin{aligned} d\langle X_t g, h \rangle_H = & (n\langle Qg, h \rangle_H + \langle A^* g, X_t h \rangle_H + \langle X_t g, A^* h \rangle_H) dt \\ & + \langle \sqrt{X_t} dW_t \sqrt{Q} g, h \rangle_H + \langle \sqrt{Q} dW_t^* \sqrt{X_t} g, h \rangle_H. \end{aligned}$$

(Wis_∞)

Theorem (C., Cuchiero, Khedher (2023))

Assume

- ▶ $A: D(A) \subseteq H \rightarrow H$ generator of a C_0 -semigroup $(e^{tA})_{t \geq 0}$ **and** e^{tA} **is injective for some $t > 0$,**
- ▶ $Q \in S^+(H)$ **is injective** and $\int_0^t \|Qe^{sA}\|_{L_2(H)} ds < \infty$ for all $t > 0$,
- ▶ $\alpha \in \mathbb{R}$,
- ▶ W is an $L_2(H)$ -cylindrical Brownian motion W ,
- ▶ $X: [0, \infty) \times \Omega \rightarrow S_1^+(H)$ such that $\forall g, h \in D(A^*), \forall t \geq 0$:

$$\begin{aligned} d\langle X_t g, h \rangle_H = & (n\langle Qg, h \rangle_H + \langle A^* g, X_t h \rangle_H + \langle X_t g, A^* h \rangle_H) dt \\ & + \langle \sqrt{X_t} dW_t \sqrt{Q} g, h \rangle_H + \langle \sqrt{Q} dW_t^* \sqrt{X_t} g, h \rangle_H. \end{aligned}$$

(Wis_∞)

Then $\alpha \in \mathbb{N}$ and $\text{rank}(X_0) \leq \alpha$. Moreover, $\text{rank}(X_t) = \alpha$ a.s. for almost all $t \geq 0$.

Proof.

Graczyk, Malecki, and Mayerhofer (2018), Letac and Massam (2018), and Mayerhofer (2019) provide a characterization of **finite dimensional Wishart processes** via its Laplace transform.

Proof.

Graczyk, Malecki, and Mayerhofer (2018), Letac and Massam (2018), and Mayerhofer (2019) provide a characterization of **finite dimensional Wishart processes** via its Laplace transform.

This ensures $\alpha \in \mathbb{N}$.

To see that $\text{rank}(X_t) = \alpha$ a.s. for almost all $t \geq 0$ we adapt the proof of CFMT (2012) (this requires an extremely technical approximation result). □

Conclusions

- ▶ Infinite-dimensional Wishart processes exist,

Conclusions

- ▶ Infinite-dimensional Wishart processes exist, **but** it seems they are necessarily of finite (fixed) rank.

Conclusions

- ▶ Infinite-dimensional Wishart processes exist, **but** it seems they are necessarily of finite (fixed) rank.
- ▶ Both the **characteristic function** and the **Laplace transform** of infinite-dimensional Wishart processes can be given explicitly.

Conclusions

- ▶ Infinite-dimensional Wishart processes exist, **but** it seems they are necessarily of finite (fixed) rank.
- ▶ Both the **characteristic function** and the **Laplace transform** of infinite-dimensional Wishart processes can be given explicitly.
- ▶ Infinite-dimensional Wishart processes are **Feller** with respect to (a minor refinement of) the weak*-topology.

Conclusions

- ▶ Infinite-dimensional Wishart processes exist, **but** it seems they are necessarily of finite (fixed) rank.
- ▶ Both the **characteristic function** and the **Laplace transform** of infinite-dimensional Wishart processes can be given explicitly.
- ▶ Infinite-dimensional Wishart processes are **Feller** with respect to (a minor refinement of) the weak*-topology.

Some advertisement: a pure jump model

A tractable covariance model that can be of infinite rank:

Some advertisement: a pure jump model

A **tractable** covariance model that **can be of infinite rank**:

C., Karbach, and Khedher (2022) establish existence of a process $X: [0, \infty) \times \Omega \rightarrow S_2^+(H)$ satisfying

$$dX_t = (b + BX_t) dt + dJ_t^X,$$

where

Some advertisement: a pure jump model

A **tractable** covariance model that **can be of infinite rank**:

C., Karbach, and Khedher (2022) establish existence of a process $X: [0, \infty) \times \Omega \rightarrow S_2^+(H)$ satisfying

$$dX_t = (b + BX_t) dt + dJ_t^X,$$

where

- ▶ $b \in S^+(H)$;
- ▶ $B \in \mathcal{L}(L^2(H))$;
- ▶ J^X a pure jump process whose compensator ν is an affine function of X ; i.e., $\nu(x, d\xi) = m(d\xi) + \langle \frac{\mu(d\xi)}{\|\xi\|^2}, x \rangle$, with $m: \mathcal{B}(S^+(H)) \rightarrow [0, \infty]$ and $\mu: \mathcal{B}(S^+(H)) \rightarrow S^+(H)$,

provided b , B , m , and μ satisfy some admissibility conditions.

Open questions and outlook

- ▶ Work in progress: equations for the eigenvalues and -vectors of infinite-dimensional Wishart processes (global existence?).

Open questions and outlook

- ▶ Work in progress: equations for the eigenvalues and -vectors of infinite-dimensional Wishart processes (global existence?).
- ▶ Are there Wishart processes that are **not of finite rank**? (e.g. if Q is not injective)?

Open questions and outlook

- ▶ Work in progress: equations for the eigenvalues and -vectors of infinite-dimensional Wishart processes (global existence?).
- ▶ Are there Wishart processes that are **not of finite rank**? (e.g. if Q is not injective)?
- ▶ Are there other 'nice' (tractable) stochastic processes taking values in $S_1^+(H)$ (preferably **not** of finite rank)?

References

- ▶ Cox, S., Cuchiero, C., and Khedher, A.: *Infinite-dimensional Wishart processes*. arXiv 2304.03490.
- ▶ Cox, S., Karbach, S., and Khedher, A.: *Affine pure-jump processes on positive Hilbert-Schmidt operators*. Stoch. Process. Appl. (2022)
- ▶ Cox, S., Karbach, S., and Khedher, A.: *An infinite-dimensional affine stochastic volatility model*. Math. Fin. (2022)