

On the Periodic Model of Uncertainty Quantification With Application to Inverse Problems

Vesa Kaarnioja (FU Berlin)

Thanks to collaborators: Yoshihito Kazashi (Uni. Strathclyde), Frances Kuo (UNSW Sydney), Fabio Nobile (EPFL), and Ian Sloan (UNSW Sydney)

Table of contents

Part I: Quasi-Monte Carlo (QMC) cubature using lattice rules

Part II: The periodic model of uncertainty quantification for PDEs

Part III: Kernel interpolation over lattice point sets

Part IV: Application to an inverse problem

Part I: Quasi-Monte Carlo cubature

Lattice rules


Rank-1 lattice rules

$$Q_{s,n}(f) = \frac{1}{n} \sum_{i=1}^n f(\mathbf{t}_i) \approx \int_{[0,1]^s} f(\mathbf{y}) \, d\mathbf{y} = I_s(f)$$

have the points

$$\mathbf{t}_i = \text{mod} \left(\frac{i\mathbf{z}}{n}, 1 \right), \quad i \in \{1, \dots, n\},$$

where the entire point set is determined by the *generating vector* $\mathbf{z} \in \mathbb{N}^s$, with all components *coprime* to n .



Lattice rule with $\mathbf{z} = (1, 55)$ and $n = 89$
nodes in $[0, 1]^2$

Lattice rules and periodic functions are a match made in heaven!

Periodic means

$$f(y_1, y_2, \dots, y_s) = f(y_1 + 1, y_2, \dots, y_s) = f(y_1, y_2 + 1, \dots, y_s) = \dots$$

Dimension $s = 1$: the only lattice is the left-Riemann rule

For $z \in \{1, \dots, n-1\}$, $\gcd(z, n) = 1$, it holds that

$$Q_{1,n}(f) = \frac{1}{n} \sum_{k=1}^n f\left(\bmod\left(\frac{kz}{n}, 1\right)\right) = \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right).$$

Suppose $f: [0, 1) \rightarrow \mathbb{R}$ is p times continuously differentiable and periodic.

Let $h = \frac{1}{n}$. Then the Euler–Maclaurin summation formula gives

$$\begin{aligned} \sum_{k=0}^{n-1} hf(kh) &= \int_0^1 f(x) \, dx + \sum_{k=1}^{\lfloor p/2 \rfloor} \frac{B_{2k}}{(2k)!} (f^{(2k-1)}(1) - f^{(2k-1)}(0)) \\ &\quad - (-1)^p h^p \int_0^1 \tilde{B}_p(x) f^{(p)}(x) \, dx \end{aligned}$$

$$\left| \int_0^1 f(x) \, dx - \frac{1}{n} \sum_{k=0}^{n-1} f\left(\frac{k}{n}\right) \right| = O(n^{-p}).$$

Dimension $s = 1$: the only lattice is the left-Riemann rule

For $z \in \{1, \dots, n-1\}$, $\gcd(z, n) = 1$, it holds that

$$Q_{1,n}(f) = \frac{1}{n} \sum_{k=1}^n f\left(\bmod\left(\frac{kz}{n}, 1\right)\right) = \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right).$$

Suppose $f: [0, 1) \rightarrow \mathbb{R}$ is p times continuously differentiable and periodic. Let $h = \frac{1}{n}$. Then the Euler–Maclaurin summation formula gives

$$\begin{aligned} \sum_{k=0}^{n-1} hf(kh) &= \int_0^1 f(x) dx + \sum_{k=1}^{\lfloor p/2 \rfloor} \frac{B_{2k}}{(2k)!} \overbrace{(f^{(2k-1)}(1) - f^{(2k-1)}(0))}^{=0} \\ &\quad - (-1)^p h^p \int_0^1 \tilde{B}_p(x) f^{(p)}(x) dx \end{aligned}$$

$$\left| \int_0^1 f(x) dx - \frac{1}{n} \sum_{k=1}^{n-1} f\left(\frac{k}{n}\right) \right| = O(n^{-p})$$

Dimension $s = 1$: the only lattice is the left-Riemann rule

For $z \in \{1, \dots, n-1\}$, $\gcd(z, n) = 1$, it holds that

$$Q_{1,n}(f) = \frac{1}{n} \sum_{k=1}^n f\left(\bmod\left(\frac{kz}{n}, 1\right)\right) = \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right).$$

Suppose $f: [0, 1) \rightarrow \mathbb{R}$ is p times continuously differentiable and periodic. Let $h = \frac{1}{n}$. Then the Euler–Maclaurin summation formula gives

$$\begin{aligned} \sum_{k=0}^{n-1} hf(kh) &= \int_0^1 f(x) \, dx + \sum_{k=1}^{\lfloor p/2 \rfloor} \frac{B_{2k}}{(2k)!} \overbrace{\left(f^{(2k-1)}(1) - f^{(2k-1)}(0)\right)}^{=0} \\ &\quad - (-1)^p h^p \int_0^1 \tilde{B}_p(x) f^{(p)}(x) \, dx \\ &= \int_0^1 f(x) \, dx + \mathcal{O}(h^p). \end{aligned}$$

$$\left| \int_0^1 f(x) \, dx - \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) \right| = \mathcal{O}(n^{-p})$$

Dimension $s = 1$: the only lattice is the left-Riemann rule

For $z \in \{1, \dots, n-1\}$, $\gcd(z, n) = 1$, it holds that

$$Q_{1,n}(f) = \frac{1}{n} \sum_{k=1}^n f\left(\bmod\left(\frac{kz}{n}, 1\right)\right) = \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right).$$


Suppose $f: [0, 1) \rightarrow \mathbb{R}$ is p times continuously differentiable and periodic. Let $h = \frac{1}{n}$. Then the Euler–Maclaurin summation formula gives

$$\begin{aligned} \sum_{k=0}^{n-1} hf(kh) &= \int_0^1 f(x) dx + \sum_{k=1}^{\lfloor p/2 \rfloor} \frac{B_{2k}}{(2k)!} \overbrace{\left(f^{(2k-1)}(1) - f^{(2k-1)}(0)\right)}^{=0} \\ &\quad - (-1)^p h^p \int_0^1 \tilde{B}_p(x) f^{(p)}(x) dx \\ &= \int_0^1 f(x) dx + \mathcal{O}(h^p). \end{aligned}$$

$$\therefore \left| \int_0^1 f(x) dx - \frac{1}{n} \sum_{k=0}^{n-1} f\left(\frac{k}{n}\right) \right| = \mathcal{O}(n^{-p}).$$


Exponential convergence for analytic, periodic functions

$$\int_0^1 \exp(-\sin(2\pi x)) dx$$




Exponential convergence for analytic, periodic functions

$$\int_0^1 \exp(-\sin(2\pi x)) dx$$




Exponential convergence for analytic, periodic functions


$$\int_0^1 \exp(-\sin(2\pi x)) dx$$




Exponential convergence for analytic, periodic functions



Exponential convergence for analytic, periodic functions





Exponential convergence for analytic, periodic functions





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?


$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$



Can we observe exponential convergence with lattice rules for analytic, periodic functions when dimension $s = 2$?

$$\int_0^1 \int_0^1 \exp(-\sin(2\pi x) \cos(2\pi y)^2) dx dy$$





A suitable generating vector for an integrand satisfying certain smoothness properties can be found using a *component-by-component* (CBC) algorithm (Nuyens and Cools 2006; Kuo, Nuyens, and Cools 2006).

- For integrands belonging to certain weighted Sobolev spaces of smooth functions, the CBC algorithm can be used to produce a generating vector satisfying a rigorous error bound. As input, the CBC algorithm takes the weights and smoothness parameter of the Sobolev space (and number of QMC nodes n).
- Fast CBC: FFT can be used to reduce the computational complexity of the CBC algorithm.

Part II: The periodic model of uncertainty quantification for PDEs

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $D \subset \mathbb{R}^d$, $d \in \{1, 2, 3\}$, a bounded physical domain with Lipschitz boundary.

Elliptic PDE with uncertain/random coefficient

Find $u: D \times \Omega \rightarrow \mathbb{R}$ that satisfies

$$\begin{aligned} -\nabla \cdot (a(\mathbf{x}, \omega) \nabla u(\mathbf{x}, \omega)) &= f(\mathbf{x}) && \text{for } \mathbf{x} \in D, \\ + \text{boundary conditions} &&& \text{on } \partial D \end{aligned}$$

for almost all events $\omega \in \Omega$. Here, the diffusion coefficient $a(\cdot, \omega) \in L_+^\infty(D)$ is *uncertain*.

In forward uncertainty quantification, one is interested in computing certain response statistics of the solution, usually $\mathbb{E}[u]$ or $\mathbb{E}[G(u)]$ and $\text{Var}[u]$ or $\text{Var}[G(u)]$, where G is a (linear) functional representing some quantity of interest derived from the solution.

Depending on the application, two common models for the random field A that appear in the literature are

- uniform and affine;
- lognormal.

Background

A popular model in the literature: the uniform and affine model

For $\mathbf{x} \in D$ and $\omega \in \Omega$,

$$a(\mathbf{x}, \omega) = \bar{a}(\mathbf{x}) + \sum_{j \geq 1} Y_j(\omega) \psi_j(\mathbf{x}), \quad Y_j \text{ i.i.d. uniform on } [-\frac{1}{2}, \frac{1}{2}].$$

Computing $\mathbb{E}[u(\mathbf{x}, \cdot)]$ (or some quantity of interest $\mathbb{E}[G(u)]$) using

- Rank-1 lattice cubature rules with random shifts
 \Rightarrow cubature error $\mathcal{O}(n^{-1+\varepsilon})$ at best. (Kuo, Schwab, Sloan 2012)
- Interlaced polynomial lattice rules
 \Rightarrow higher order convergence $\mathcal{O}(n^{-1/p})$ for some $0 < p < 1$ (p is a summability exponent s.t. $(\|\psi_j\|_{L^\infty})_{j \geq 1} \in \ell^p$). (Dick, Kuo, Le Gia, Nuyens, Schwab 2014)

Periodic model of UQ

In this talk, we instead model the uncertainty in the diffusion coefficient as follows.

For $\mathbf{x} \in D$ and $\omega \in \Omega$,

$$a(\mathbf{x}, \omega) = \bar{a}(\mathbf{x}) + \sum_{j \geq 1} \Theta(Y_j(\omega)) \psi_j(\mathbf{x}), \quad Y_j \text{ i.i.d. uniform on } [-\frac{1}{2}, \frac{1}{2}]$$


with the special choice $\Theta(y) = \sin(2\pi y)$.

- Note that $Z(\omega) := \sin(2\pi Y(\omega))$ has the probability density $\frac{1}{\pi} \frac{1}{\sqrt{1-z^2}}$ on $[-1, 1]$, i.e. $Z \sim \text{Arcsine}(-1, 1)$.
- We can match the mean and covariance of a with the “uniform model” by choosing $\Theta(y) = \frac{1}{\sqrt{6}} \sin(2\pi y)$.
- Note that the periodicity is only assumed for the *random/uncertain variable*!

Affine vs. periodic


Affine

$$a(x, \mathbf{y}) = \bar{a}(x) + \sum_{j=1}^{100} y_j \psi_j(x)$$



Periodic

$$a(x, \mathbf{y}) = \bar{a}(x) + \frac{1}{\sqrt{6}} \sum_{j=1}^{100} \sin(2\pi y_j) \psi_j(x)$$



$$\bar{a}(x) = 2, \quad \psi_j(x) = j^{-3/2} \sin((j - \frac{1}{2})\pi x), \quad x \in [0, 1]$$

Let $U := [-1/2, 1/2]^{\mathbb{N}}$ and $D \subset \mathbb{R}^d$, $d \in \{1, 2, 3\}$, a nonempty bounded Lipschitz domain. For the parametric PDE

$$\begin{cases} -\nabla \cdot (a(\mathbf{x}, \mathbf{y}) \nabla u(\mathbf{x}, \mathbf{y})) = f(\mathbf{x}) & \text{for } \mathbf{x} \in D, \mathbf{y} \in U \\ u(\mathbf{x}, \mathbf{y}) = 0 & \text{for } \mathbf{x} \in \partial D, \mathbf{y} \in U, \end{cases}$$

with $u(\cdot, \mathbf{y}) \in H_0^1(D)$, $f \in H^{-1}(D)$, and

$$a(\mathbf{x}, \mathbf{y}) = \bar{a}(\mathbf{x}) + \sum_{j=1}^s \sin(2\pi y_j) \psi_j(\mathbf{x}),$$

with assumptions

- $0 < a_{\min} \leq a(\mathbf{x}, \mathbf{y}) \leq a_{\max} < \infty$ for all $\mathbf{x} \in D$, $\mathbf{y} \in U$
- $\sum_{j=1}^{\infty} \|\psi_j\|_{L^\infty}^p < \infty$ for some $p \in (0, 1)$
- $\|\psi_1\|_{L^\infty} \geq \|\psi_2\|_{L^\infty} \geq \dots$

[K–Kuo–Sloan 2020] showed that there exists a constructible lattice rule satisfying the QMC cubature error

$$|I_s(G(u)) - Q_{n,s}(G(u))| \leq C n^{-1/p} \quad \text{with constant } C > 0 \text{ independent of } s,$$

for any linear quantity of interest $G: H_0^1(D) \rightarrow \mathbb{R}$.

Numerical example: QMC for PDE [K–Kuo–Sloan (2020)]

Let us consider the PDE problem

$$-\nabla \cdot (a_{\text{per}}(\mathbf{x}, \mathbf{y}) \nabla u(\mathbf{x}, \mathbf{y})) = x_2, \quad u(\cdot, \mathbf{y})|_{\partial D} = 0,$$

in the physical domain $D = (0, 1)^2$ with the diffusion coefficient

$$a_{\text{per}}(\mathbf{x}, \mathbf{y}) = 2 + \sum_{j=1}^{100} \sin(2\pi y_j) \psi_j(x), \quad y_j \in [-\tfrac{1}{2}, \tfrac{1}{2}],$$

where $\psi_j(\mathbf{x}) = \frac{c}{\sqrt{6}} j^{-\theta} \sin(j\pi x_1) \sin(j\pi x_2)$. **Note that** $\|\psi_j\|_{L^\infty} \propto j^{-\theta}$.




Figure 7: Left: $\theta = 2$. Right: $\theta = 4$.

Part III: Kernel interpolation over lattice point sets

Let us continue the study of our elliptic model PDE problem.

In [K–Kazashi–Kuo–Nobile–Sloan (2022)], we studied *kernel interpolation of smooth, periodic functions based on lattice point sets*. We considered the following setting:

- Let $\alpha \geq 1$ be an integer and let $H := H_{s,\alpha,\gamma}$ be the Hilbert space containing absolutely continuous, somewhat smooth periodic functions $f: [0, 1]^s \rightarrow \mathbb{R}$ endowed with the norm

$$\|f\|_H^2 := \sum_{u \subseteq \{1:s\}} \frac{1}{(2\pi)^{2\alpha|u|}\gamma_u} \int_{[0,1]^{|u|}} \left| \int_{[0,1]^{s-|u|}} \left(\prod_{j \in u} \frac{\partial^\alpha}{\partial y_j^\alpha} \right) f(\mathbf{y}) \, d\mathbf{y}_{-u} \right|^2 d\mathbf{y}_u,$$

provided that f has mixed partial derivatives of order α .

The space H is actually a *reproducing kernel Hilbert space* (RKHS), with an explicitly known and analytically simple reproducing kernel:

$$K(\mathbf{y}, \mathbf{y}') := \sum_{\mathbf{u} \subseteq \{1:s\}} \gamma_{\mathbf{u}} \prod_{j \in \mathbf{u}} \eta_{\alpha}(y_j, y'_j),$$

where

$$\eta_{\alpha}(y, y') = \frac{(2\pi)^{2\alpha}}{(-1)^{\alpha+1}(2\alpha)!} B_{2\alpha}(\text{frac}(y - y')), \quad y, y' \in [0, 1],$$

where $B_2(y) = y^2 - y + \frac{1}{6}$, $B_4(y) = y^4 - 2y^3 + y^2 - \frac{1}{30}$, and so on, are the *Bernoulli polynomials*. In particular,

$$\langle f, K(\cdot, \mathbf{y}) \rangle_H = f(\mathbf{y}) \quad \text{for all } f \in H \text{ and } \mathbf{y} \in [0, 1]^s.$$

Example: If $(\gamma_{\mathbf{u}})_{\mathbf{u} \subseteq \{1,\dots,s\}}$ are *product weights*, i.e.,

$$\gamma_{\mathbf{u}} := \prod_{j \in \mathbf{u}} \gamma_j, \quad \mathbf{u} \subseteq \{1, \dots, s\},$$

then

$$K(\mathbf{y}, \mathbf{y}') = \prod_{j=1}^s (1 + \gamma_j \eta_{\alpha}(y_j, y'_j)).$$

Suppose that one is interested in finding an approximation for the function $f \in H$ based on the point evaluations $f(\mathbf{t}_1), \dots, f(\mathbf{t}_n)$, $\mathbf{t}_j \in [0, 1]^s$. We introduce the *kernel interpolant*

$$f_n(\mathbf{y}) := \sum_{k=1}^n c_k K(\mathbf{t}_k, \mathbf{y}), \quad \mathbf{t}_k := \text{mod}\left(\frac{k\mathbf{z}}{n}, 1\right), \quad (1)$$

and require the interpolation property $f_n(\mathbf{t}_k) = f(\mathbf{t}_k)$ for hold for all $k = 1, \dots, n$. Then the coefficients can be solved from the linear system

$$\mathbf{K}\mathbf{c} = \mathbf{f},$$

where $\mathbf{c} := [c_1, \dots, c_n]^T$ are the coefficients in (1) and

$$K_{k,\ell} = K(\mathbf{t}_k, \mathbf{t}_\ell) \quad \text{and} \quad \mathbf{f} := [f(\mathbf{t}_1), \dots, f(\mathbf{t}_n)]^T.$$

Note that $K_{k,\ell} = K(\frac{(k-\ell)\mathbf{z}}{n}, \mathbf{0})$, i.e., \mathbf{K} is a *circulant matrix* \Rightarrow

$$\mathbf{c} = \text{ifft}(\text{fft}(\mathbf{f})./\text{fft}(\mathbf{K}_{:,1}))$$

This can be computed in $\mathcal{O}(n \log n)$ time!

The kernel interpolant is cheap to construct!

In analogy to the cubature setting, the PDE problem

$$\begin{cases} -\nabla \cdot (a(\mathbf{x}, \mathbf{y}) \nabla u(\mathbf{x}, \mathbf{y})) = f(\mathbf{x}) & \text{for } \mathbf{x} \in D, \mathbf{y} \in U, \\ u(\mathbf{x}, \mathbf{y}) = 0 & \text{for } \mathbf{x} \in \partial D, \mathbf{y} \in U, \end{cases}$$

with $u(\cdot, \mathbf{y}) \in H_0^1(D)$, $f \in H^{-1}(D)$, and

$$a(\mathbf{x}, \mathbf{y}) = \bar{a}(\mathbf{x}) + \sum_{j=1}^s \sin(2\pi y_j) \psi_j(\mathbf{x})$$

and assumptions

- $0 < a_{\min} \leq a(\mathbf{x}, \mathbf{y}) \leq a_{\max} < \infty$ for all $\mathbf{x} \in D, \mathbf{y} \in U$
- $\sum_{j=1}^{\infty} \|\psi_j\|_{L^\infty}^p < \infty$ for some $p \in (0, 1)$
- $\|\psi_1\|_{L^\infty} \geq \|\psi_2\|_{L^\infty} \geq \dots$

[K–Kazashi–Kuo–Nobile–Sloan 2022] showed that there exists a sequence of SPOD weights (entering both the expression of the kernel K in the interpolant and as inputs to a CBC algorithm) and a constructible lattice rule satisfying the kernel approximation error

$$\|u - u_n\|_{L^2(U \times D)} = \mathcal{O}(n^{-\frac{1}{2p} + \frac{1}{4}}) \quad \text{with constant } C > 0 \text{ independent of } s.$$

Kernel approximation for PDE: L^2 error



Let us consider the PDE problem

$$-\nabla \cdot (a_{\text{per}}(\mathbf{x}, \mathbf{y}) \nabla u(\mathbf{x}, \mathbf{y})) = x_2, \quad u(\cdot, \mathbf{y})|_{\partial D} = 0,$$

in the physical domain $D = (0, 1)^2$ with the diffusion coefficient

$$a_{\text{per}}(\mathbf{x}, \mathbf{y}) = 1 + \sum_{j=1}^{100} \sin(2\pi y_j) \psi_j(x), \quad y_j \in [0, 1],$$

where $\psi_j(\mathbf{x}) = c j^{-\theta} \sin(j\pi x_1) \sin(j\pi x_2)$. **Note that $\|\psi_j\|_{L^\infty} \propto j^{-\theta}$.**



Reducing the computational complexity

The SPOD weights used in the construction of the kernel interpolant were

$$\gamma_u := \sum_{\mathbf{m}_u \in \{1:\alpha\}^{|\mathbf{u}|}} (|\mathbf{m}_u|!)^{\frac{2}{1+\lambda}} \prod_{j \in \mathbf{u}} \left(\frac{b_j^{m_j} S(\alpha, m_j)}{\sqrt{2e^{1/e} \zeta(2\alpha\lambda)}} \right)^{\frac{2}{1+\lambda}}. \quad (2)$$

- the cost to obtain the generating vector \mathbf{z} is $\mathcal{O}(s n \log n + s^3 \alpha^2 n)$;
- the cost of evaluating the kernel interpolant is $\mathcal{O}(s^2 \alpha^2 n)$.

New idea (see Ian's talk on Friday): leave out the order-dependent part $(|\mathbf{m}_u|!)^{\frac{2}{1+\lambda}}$ in (2), get

$$\tilde{\gamma}_u := \sum_{\mathbf{m}_u \in \{1:\alpha\}^{|\mathbf{u}|}} \prod_{j \in \mathbf{u}} \left(\frac{b_j^{m_j} S(\alpha, m_j)}{\sqrt{2e^{1/e} \zeta(2\alpha\lambda)}} \right)^{\frac{2}{1+\lambda}} = \prod_{j \in \mathbf{u}} \left(\sum_{m=1}^{\alpha} \left(\frac{b_j^m S(\alpha, m)}{\sqrt{2e^{1/e} \zeta(2\alpha\lambda)}} \right)^{\frac{2}{1+\lambda}} \right).$$

These are *product weights* (**“serendipitous weights”**), where

- the cost to obtain the generating vector \mathbf{z} is $\mathcal{O}(s n \log n)$;
- the cost of evaluating the kernel interpolant is $\mathcal{O}(s n)$.

Kernel approximation for PDE: L^2 error (redux)



Let us consider the PDE problem

$$-\nabla \cdot (a_{\text{per}}(\mathbf{x}, \mathbf{y}) \nabla u(\mathbf{x}, \mathbf{y})) = x_2, \quad u(\cdot, \mathbf{y})|_{\partial D} = 0,$$



in the physical domain $D = (0, 1)^2$ with the diffusion coefficient

$$a_{\text{per}}(\mathbf{x}, \mathbf{y}) = 1 + \sum_{j=1}^{100} \sin(2\pi y_j) \psi_j(x), \quad y_j \in [0, 1],$$




where $\psi_j(\mathbf{x}) = c j^{-\theta} \sin(j\pi x_1) \sin(j\pi x_2)$. **Note that $\|\psi_j\|_{L^\infty} \propto j^{-\theta}$.**



In certain situations, the product weights can outperform SPOD weights.



The product weights can be used to perform computations for higher dimensional problems (here, $s = 1000$).



Part IV: Application to an inverse problem


The complete electrode model

Let $D := \{\mathbf{x} \in \mathbb{R}^2 : \|\mathbf{x}\| \leq 1\}$. Let $\{E_k\}_{k=1}^L \subseteq \partial D$ be an array of $L := 16$ equidistantly spaced non-overlapping electrodes of width 0.2 on the boundary ∂D . Fix the current feed $\mathbf{I} \in \mathbb{R}_{\diamond}^L$ and let $\sigma \in L_+^\infty(D)$. The *forward problem* is to find the electromagnetic potential $u \in H^1(D)$ as well as $\mathbf{U} \in \mathbb{R}^L$, the potentials on the electrodes, which satisfy

$$\begin{cases} \nabla \cdot (\sigma \nabla u) = 0 & \text{in } D, \\ \sigma \frac{\partial u}{\partial \mathbf{n}} = 0 & \text{on } \partial D \setminus \bigcup_{k=1}^L \overline{E_k}, \\ u + z_k \sigma \frac{\partial u}{\partial \mathbf{n}} = U_k & \text{on } E_k, \ k \in \{1, \dots, L\}, \\ \int_{E_k} \sigma \frac{\partial u}{\partial \mathbf{n}} \, dS = I_k, & k \in \{1, \dots, L\}, \end{cases}$$

with \mathbf{n} denoting the outer normal.

Moreover, we take $z_k = 1 \ \forall k$.



The forward problem is solved numerically using EIDORS software (FEM).

Fix the current pattern $\mathbf{l}_k := \mathbf{e}_1 - \mathbf{e}_{k+1} \in \mathbb{R}_{\diamond}^L$, $k \in \{1, \dots, L-1\}$.

Kernel-based surrogate for the forward problem: Let us parameterize the conductivity as

$$\sigma(\mathbf{x}, \mathbf{y}) := 1 + \frac{1}{\sqrt{6}} \sum_{k=1}^s \sin(2\pi y_k) \psi_k(\mathbf{x}), \quad \mathbf{x} \in D, \quad \mathbf{y} \in [0, 1]^s,$$

where $\psi_k(\mathbf{x}) := \frac{1}{(i_k^2 + j_k^2)^{\vartheta}} \sin(\pi i_k \frac{x_1+1}{2}) \sin(\pi j_k \frac{x_2+1}{2})$, $\vartheta = 1.2$, the sequence $(i_k, j_k)_{k \geq 1}$ is an ordering of the elements of $\mathbb{N} \times \mathbb{N}$ s.t. $\|\psi_k\|_{L^\infty} = \mathcal{O}(k^{-\vartheta})$ by Weyl's asymptotics. We set $s = 30$.

Denote by $\mathcal{U}(\mathbf{y}) := \text{vec}([\mathbf{U}_1, \dots, \mathbf{U}_{L-1}]) \in \mathbb{R}^{L(L-1)}$ the (flattened) voltage matrix, comprised of the electrode potential measurements corresponding to the current pattern $\mathbf{l}_1, \dots, \mathbf{l}_{L-1}$ and $\mathbf{y} \in [0, 1]^s$.

We construct the (vector-valued) QMC-kernel interpolant $\mathcal{U}_n(\mathbf{y}) := \sum_{k=1}^n \mathbf{c}_k K(\mathbf{t}_k, \mathbf{y}) \in \mathbb{R}^{L(L-1)}$ (using **serendipitous weights**) for the mapping $G: \mathbf{y} \mapsto \mathcal{U}(\mathbf{y})$ based on $n = 1\,024\,207$ QMC nodes satisfying $G(\mathbf{t}_k) = \mathcal{U}_n(\mathbf{t}_k) \forall k$.

Experiment setup

We have constructed the QMC–kernel interpolant $\mathcal{U}_n(\mathbf{y})$ offline based on the periodically parameterized model for $\sigma(\mathbf{x}, \mathbf{y})$. For the numerical experiments, we

- fix some target conductivity σ_{target} and numerically compute the “exact” electrode potential measurements $\mathbf{U}_{\text{exact}} := \text{vec}([\mathbf{U}_{\text{exact}}^1, \dots, \mathbf{U}_{\text{exact}}^{L-1}]) \in \mathbb{R}^{L(L-1)}$. To avoid the inverse crime, we do not use the same FE mesh that was used to build the surrogate; instead we use a finer FE mesh.
- we contaminate the electrode potential measurements with noise

$$\mathbf{U}_{\text{noisy}} = \mathbf{U}_{\text{exact}} + \boldsymbol{\eta}, \quad \boldsymbol{\eta} \sim \mathcal{N}(\mathbf{0}, \tau^2 \mathbf{I}),$$

$$\text{where } \tau := 10^{-3} \max_{j,k=1,\dots,L(L-1)} |(U_{\text{exact}})_j - (U_{\text{exact}})_k|.$$

Our reconstruction is $\sigma(\mathbf{x}, \mathbf{y}^*)$, where

$$\mathbf{y}^* := \arg \min_{\mathbf{y} \in [0,1]^s} \{\|\mathbf{U}_{\text{noisy}} - \mathcal{U}_n(\mathbf{y})\|^2\}.$$

The minimization is carried out using `lsqnonlin` in MATLAB with the levenberg-marquardt algorithm.




Figure 10: Left: target conductivity. Right: reconstructed conductivity.




Figure 11: Left: target conductivity. Right: reconstructed conductivity.

Conclusions

- Kernel interpolation method that can be used to approximate the output high-dimensional parametric PDEs. Kernel interpolant can be constructed efficiently at cost $\mathcal{O}(n \log n)$. No multi-index sets! (Compare with sparse grids or trigonometric approximation.)
- Using product weights, practical for challenging high-dimensional problems (e.g., as surrogates for Bayesian inversion).
- For EIT, the kernel interpolation scheme could be useful for efficient recovery of other uncertainties (domain shape, electrode positions, contact resistances, etc.).

References



K., F. Y. Kuo, and I. H. Sloan. Uncertainty quantification using periodic random variables. *SIAM J. Numer. Anal.* **58**(2):1068–1091, 2020.



K., Y. Kazashi, F. Y. Kuo, F. Nobile, and I. H. Sloan. Fast approximation by periodic kernel-based lattice-point interpolation with application in uncertainty quantification. *Numer. Math.* **150**:33–77, 2022.



K., F. Y. Kuo, and I. H. Sloan. Lattice-based kernel approximation and serendipitous weights for parametric PDEs in very high dimensions. Preprint arXiv:2303.17755, 2023.